

**A COMBINED GROWTH FACTOR-DELETED AND THYMIDINE KINASE-  
DELETED VACCINIA VIRUS VECTOR**

Related Applications

sub  
A1

[0001] This application claims the benefit of priority from PCT/US00/14679, filed May 26, 2000, which claims the benefit of priority from U.S. Provisional Patent Application No. 60/137,126, filed May 28, 1999, each of which is hereby incorporated by reference in their entirety.

Background of the Invention

Field of the Invention

FOETT-TELEFON

[0002] The present invention relates to mutant vaccinia virus expression vectors. The mutant expression vectors of the present invention show substantially no virus replication in non-dividing cells and as such are superior to previous vaccinia virus expression vectors. More specifically, the vaccinia virus expression vectors of the present invention are associated with negative thymidine kinase and vaccinia virus growth factor phenotypes. The vaccinia virus vectors of the present invention are suitable for use as vaccines, cancer therapies, as well as for gene delivery vectors.

Description of the Related Art

[0003] Vaccinia virus has a long history of service as an expression vehicle. Vaccinia virus has been used most notably as a vaccination against smallpox, however, active immunization against smallpox is far less common since the eradication of the disease in 1977. Today, as a result of various advances in molecular biology, the vaccinia virus show promise as a vaccine against diseases other than smallpox.

[0004] The vaccinia virus genome can be manipulated using general molecular biology techniques well known in the art, to perform as an expression vector for exogenous, non-vaccinia virus genes. In one study, a region of the wild-type (WT) strain of vaccinia virus was shown to be suitable for the insertion and expression of foreign DNA. Mackett, et al., *Proc Natl Acad Sci USA*, 79(23):7415-9 (1982). The ability to express foreign DNA by

Altitude	Temperature	Humidity	Wind	Clouds	Pressure	Time	Remarks
1000	15.0	85	10	10	1013.2	0800	Clear
1000	14.5	85	10	10	1013.2	0815	Clear
1000	14.0	85	10	10	1013.2	0830	Clear
1000	13.5	85	10	10	1013.2	0845	Clear
1000	13.0	85	10	10	1013.2	0900	Clear
1000	12.5	85	10	10	1013.2	0915	Clear
1000	12.0	85	10	10	1013.2	0930	Clear
1000	11.5	85	10	10	1013.2	0945	Clear
1000	11.0	85	10	10	1013.2	1000	Clear
1000	10.5	85	10	10	1013.2	1015	Clear
1000	10.0	85	10	10	1013.2	1030	Clear
1000	9.5	85	10	10	1013.2	1045	Clear
1000	9.0	85	10	10	1013.2	1100	Clear
1000	8.5	85	10	10	1013.2	1115	Clear
1000	8.0	85	10	10	1013.2	1130	Clear
1000	7.5	85	10	10	1013.2	1145	Clear
1000	7.0	85	10	10	1013.2	1200	Clear
1000	6.5	85	10	10	1013.2	1215	Clear
1000	6.0	85	10	10	1013.2	1230	Clear
1000	5.5	85	10	10	1013.2	1245	Clear
1000	5.0	85	10	10	1013.2	1300	Clear
1000	4.5	85	10	10	1013.2	1315	Clear
1000	4.0	85	10	10	1013.2	1330	Clear
1000	3.5	85	10	10	1013.2	1345	Clear
1000	3.0	85	10	10	1013.2	1400	Clear
1000	2.5	85	10	10	1013.2	1415	Clear
1000	2.0	85	10	10	1013.2	1430	Clear
1000	1.5	85	10	10	1013.2	1445	Clear
1000	1.0	85	10	10	1013.2	1500	Clear
1000	0.5	85	10	10	1013.2	1515	Clear
1000	0.0	85	10	10	1013.2	1530	Clear
1000	-0.5	85	10	10	1013.2	1545	Clear
1000	-1.0	85	10	10	1013.2	1600	Clear
1000	-1.5	85	10	10	1013.2	1615	Clear
1000	-2.0	85	10	10	1013.2	1630	Clear
1000	-2.5	85	10	10	1013.2	1645	Clear
1000	-3.0	85	10	10	1013.2	1700	Clear
1000	-3.5	85	10	10	1013.2	1715	Clear
1000	-4.0	85	10	10	1013.2	1730	Clear
1000	-4.5	85	10	10	1013.2	1745	Clear
1000	-5.0	85	10	10	1013.2	1800	Clear
1000	-5.5	85	10	10	1013.2	1815	Clear
1000	-6.0	85	10	10	1013.2	1830	Clear
1000	-6.5	85	10	10	1013.2	1845	Clear
1000	-7.0	85	10	10	1013.2	1900	Clear
1000	-7.5	85	10	10	1013.2	1915	Clear
1000	-8.0	85	10	10	1013.2	1930	Clear
1000	-8.5	85	10	10	1013.2	1945	Clear
1000	-9.0	85	10	10	1013.2	2000	Clear
1000	-9.5	85	10	10	1013.2	2015	Clear
1000	-10.0	85	10	10	1013		

**[0005]**

**[0006]**

**[0007]**

**[0008]**

DNA led to attenuation, Buller, et al., examined several TK- recombinants for their ability to replicate. Buller, et al., *Nature* 317:813-815 (1985). Although reduced viral mediated pathogenesis was observed with these mutants, the TK- viruses retained the ability to replicate in non-dividing cells.

[0009] Although the vaccinia virus has been the subject of considerable research to date, there remains a need for a vaccinia virus expression vector that substantially fails to replicate in non-dividing cells.

#### Brief Description of the Drawings

[0010] FIGURE 1 depicts viral expansion in NIH3T3 cells of wild-type (F13), VGF- phenotype vaccinia virus VVVGf- (VSC20), TK- phenotype vaccinia virus VVTK- (VJS6), and the VGF-/TK- double mutant vaccinia virus VVDD. FIGURE 1A depicts viral expansion in non-confluent NIH3T3 cells. FIGURE 1B depicts viral expansion in confluent NIH3T3 cells.

[0011] FIGURE 2 depicts the proportion of surviving mice challenged with  $10^7$  infectious particles (IP) of wild-type (F13), VGF- phenotype vaccinia virus VVVGf- (VSC20), TK- phenotype vaccinia virus VVTK- (VJS6), and the VGF-/TK- double mutant vaccinia virus VVKK.

[0012] FIGURE 3 depicts the proportion of surviving mice challenged with  $10^8$  IP of wild-type (F13), VGF- phenotype vaccinia virus VVVGf- (VSC20), TK- phenotype vaccinia virus VVTK- (VJ86), and the VGF-/TK- double mutant vaccinia virus VVDD.

[0013] FIGURE 4 depicts the proportion of surviving mice challenged with  $10^8$  IP of wild-type (F13), VGF- phenotype vaccinia virus (VSC20) VVVGf-, TK- phenotype vaccinia virus VVTK- (VJS6), and the VGF-/TK- double mutant vaccinia virus VVDDEGFP, which contains the gene for the green fluorescent protein.

[0014] FIGURE 5 depicts median viral titers sampled from brain and tumor tissue after administration of  $10^7$  pfu IP to tumor bearing nude mice. Viral titers are shown as pfu/mg protein.

[0015] FIGURE 6 depicts changes in mean tumor volume after administration of  $10^9$  pfu IP of the double mutant VVDDEGFP and virus free buffer. Change in tumor volume is shown in cubic millimeters ( $\text{mm}^3$ ).

#### Detailed Description of the Preferred Embodiment

[0016] The present invention relates to poxvirus expression vectors that are associated with TK and VGF negative phenotypes. The vectors of the present invention are substantially incapable of replicating in non-dividing cells. Substantially incapable of replication means a vaccinia virus that replicates 90% as well as the wild-type virus, or 80% as well, or 70% as well, or 60% as well, or 50% as well, or 40% as well, or 30% as well, or 20% as well, or 10% as well, or 1% as well, or any percent within this described field. More specifically, the present invention relates to vaccinia virus expression vectors that are associated with TK and VGF negative phenotypes. The present invention also relates to methods of making the vaccinia virus discussed herein, as well as to methods of using these expression vectors.

[0017] The expression vectors of the present invention are constructed from members of *Poxviridae* that possess a thymidine kinase phenotype and a virus associated growth factor phenotype. Expression vectors from the subfamilies *Chordopoxvirinae* and *Entomopoxvirinae* are encompassed by the present invention. Specifically, virus that are members of the genera *Orthopoxvirus*, *Parapoxvirus*, *Avipoxvirus*, *Capripoxvirus*, *Leporipoxvirus*, *Suipoxvirus*, *Molluscipoxvirus*, *Yatapoxvirus*, *Entomopoxvirus A*, *Entomopoxvirus B*, and *Entomopoxvirus C* are contemplated for use with the methods of the present invention. Specific examples of viruses that may be used with the methods of the present invention include: camelpox, cowpox, ectromelia, monkeypox, racoonpox, skunkpox, taterapox, Uasin Gishu, variola, volepox, Auzdyk disease, chamois contagious ecthyma, orf, pseudocowpox, parapox of deer, sealpox, carnarypox, fowlpox, juncopox, mynahpox, pigeonpox, psittachinepox, quailpox, peacockpox, penguinpox, sparrowpox, quailpox, sparrowpox, starlingpox, turkeypox, goatpox, lumpy skin disease, sheeppox, hare fibroma, myxoma, rabbit fibroma, squirrel fibroma, swinepox, molluscum contagiosum, tanapox,

Yaba monkey tumor, *Melolontha melolontha*, *Amesacta moorei*, and *Chironimus luridus*. In a preferred embodiment, vaccinia virus is used as the expression vector.

[0018] There are a number of vaccinia virus strains presently known. Each of these strains can be used with the methods of the present invention. These strains include the WR strain (ATCC VR-119), the Wyeth strain (ATCC VR-325), the Lederle-Chorioallantoic strain (ATCC VR-325), the CL strain (ATCC VR-117), and others; all of these strains are available from the American Type Culture Collection (Manassas, VA).

[0019] Vaccinia virus was introduced in 1982 as a vector for transient expression of genes in mammalian cells (Mackett et al., 1982). As a vector, vaccinia virus has a number of useful characteristics, including a capability that permits cloning large fragments of foreign DNA (>20 kbp) with retention of infectivity, a wide host range, a relatively high level of protein synthesis, and transport, secretion, processing, and posttranslational modifications as dictated by the primary structure of the expressed protein and the cell type used. For example, *N*- and *O*-glycosylation, phosphorylation, myristylation, and cleavage, as well as assembly of expressed proteins, occur in an apparently faithful manner.

[0020] Applications of vaccinia virus expression vectors are numerous. In *in vitro* applications, recombinant vaccinia virus expression vectors have utility in the production of biologically active proteins in tissue culture, analysis of mutant forms of proteins, and determination of transport and processing signals. In addition, recombinant vaccinia viruses have been important for immunological studies. (Bennink and Yewdell, *Curr Top Microbiol Immunol* 163:153-84 (1990)). In *in vivo* applications, recombinant vaccinia virus expression vectors have utility as vaccines and gene delivery vectors for human and veterinary uses.

[0021] Although the vaccinia virus has a number of potential uses, the virus also has a number of limitations that reduce its utility. For example, the virus itself is a human pathogen that can cause death in certain immune compromised individuals with defective cell-mediated immunity. Malaise, pustule formation, progressive vaccinia, and encephalitis are also known complications of vaccinia virus infection. Furthermore, accidental infection of individuals, especially children, who come into contact with an individual infected with the vaccinia virus is also a significant complication. This phenomenon is known as eczema

vaccinatum, which occurs in eczematous subjects. Large areas of skin can be infected. Eczema vaccinatum is rarely fatal, but all cases require treatment. In view of these limitations, a vaccinia virus vector with a diminished capability to replicate in non-dividing cells would be extremely useful.

[0022] The present invention describes mutant vaccinia viruses that are substantially incapable of replicating in non-dividing cells. This inability arises from a combination of mutations in viral genes that result in a mutant virus that is substantially incapable of replication in non-dividing cells. This reduced capability can be achieved by mutating various viral genes including the ribonucleotide reductase-large subunit, the ribonucleotide reductase-small subunit, thymidylate kinase, DNA ligase, dUTPase, the thymidine kinase (TK) gene, and the vaccinia virus growth factor (VGF) gene. In a preferred embodiment, double mutations in the TK and VGF genes result in a virus with a TK- and VGF- phenotype.

[0023] It is one of the surprising observations of the present invention that the combination of TK and VGF minus phenotypes yield a vaccinia virus with a diminished ability to replicate in non-dividing cells, especially *in vivo*. That is, it replicates selectively in dividing cells. The teachings described below instruct how to make and use the mutant vaccinia viruses of the present invention.

[0024] The vaccinia virus TK gene is present in one copy in the vaccinia virus genome. To create a vaccinia virus with a TK- phenotype, it is necessary to disrupt the nucleic acid sequence of the TK gene in such a way that the TK gene no longer encodes an active and functional copy of the TK gene. Disruption of the TK gene can be achieved through an insertion, an amino acid sequence altering substitution of the existing nucleic acid sequence, or a deletion in the nucleic acid sequence of the gene. Alternatively, the entire gene sequence can be removed from the vaccinia virus genome so that all or most of the TK gene sequence is absent from the resulting recombinant vaccinia virus genome. The molecular biology techniques required to facilitate these changes are found, for example, in Current Techniques in Molecular Biology, (Ed. Ausubel, et al.) (1998).

[0025] A vaccinia virus with a negative VGF phenotype can be made in a manner similar to that described for the TK gene. For example, one or both of the VGF genes can be



[0028] An assay that is capable of determining whether or not a functional TK or VGF protein is produced by a putatively recombinant vaccinia virus can be used with the methods of the present invention. For example, to determine the presence or absence of a VGF- phenotype, a putatively recombinant vaccinia virus is produced according to the methods of the present invention and tested for its ability to produce a functional VGF protein. Such assays include determining the presence of the VGF protein by examining VGF biological activity, by immunological techniques, or by molecular biology techniques such as PCR and gene sequencing to determine the presence of a gene and its sequence, and other methods well known in the art. For example, VGF protein activity can be measured by performing EGF-receptor competition binding assays, by measuring the amount of tyrosine phosphorylation of EGF receptors as a result of VGF binding, or by observing changes in signal transduction following binding to EGF receptors. The presence or absence of TK- phenotype is assayed for by examining TK enzymatic activity, by immunological techniques, by molecular biology techniques, or other methods well known in the art. For example, cells infected with a putative TK- vaccinia virus are capable of growth in the presence of 5-bromodeoxyuridine (BrdU), a nucleotide analog that incorporates into DNA in the presence of a functional TK gene. In contrast, cells infected with TK+ phenotype virus are not capable of growth in the presence of BrdU.

[0029] Following construction of the double mutant vaccinia virus vector, a gene of interest can be placed under the control of a vaccinia virus promoter and integrated into the genome of the mutant vaccinia virus. Alternatively, expression can be achieved by transfecting a shuttle vector or plasmid containing the vaccinia promoter-controlled gene into a cell that has been infected with vaccinia virus and introducing the exogenous sequence by homologous recombination. In still another variation, the bacteriophage T7 RNA polymerase gene can be integrated into the genome of vaccinia so that a gene controlled by a T7 promoter, either in a transfected plasmid or a recombinant vaccinia virus, is expressed.

[0030] There exist a number of vaccinia virus shuttle vectors that contain vaccinia virus promoters. These shuttle vectors are appropriate means to introduce by homologous recombination exogenous nucleotide sequences into the vaccinia virus genome for expression purposes. Such nucleotide sequences can also be introduced to "knock out" or inactivate the



TK and/or VGF sequences. A table of such vectors is found in TABLE 1, which was taken from Current Techniques in Molecular Biology, (Ed. Ausubel, et al.) Unit 16.17.4 (1998).

TABLE 1

Vaccinia Virus Transfer Vectors

Vector <sup>a</sup>	Promoter <sup>b</sup>	Cloning sites <sup>c</sup>	Insertion sites <sup>d</sup>	Selection/screening	Reference <sup>f</sup>
PGS20	<i>p7.5</i> (E/L)	<i>Bam</i> HI; <i>Sma</i> I	TK	TK-	Mackett et al., 1984
PSC11	<i>p7.5</i> (E/L)	<i>Sma</i> I; MCS	TK	TK-, $\beta$ -gal	Chakrabarti et al., 1985; Earl et al., 1990; Bacik et al., 1994
PMJ601, pMJ602	<i>p</i> <sub>syn</sub> (L)	MCS	TK	TK-, $\beta$ -gal	Davison and Moss, 1990
PRB21	<i>p</i> <sub>syn</sub> (E/L)	MCS	F12L/F13L	Plaque	Blasco and Moss, 1995
PMC02	<i>p</i> <sub>syn</sub> (E/L)	MCS	TK	TK-,GUS	Carroll and Moss, 1995
PSC59	<i>p</i> <sub>syn</sub> (E/L)	MCS	TK	TK-	Chakrabarti et al., 1997
PSC65	<i>p</i> <sub>syn</sub> (E/L)	MCS	TK	TK-, $\beta$ -gal	Chakrabarti et al., 1997
PJS4	<i>p</i> <sub>syn</sub> (E/L) x2	MCS	TK	TK-	Chakrabarti et al., 1997
PJS5	<i>p</i> <sub>syn</sub> (E/L)x2	MCS	TK	TK-,gpt	Chakrabarti et al., 1997
PG06	<i>p</i> <sub>syn</sub> (E/L)x2	MCS	Del III	Transient gpt <sup>e</sup>	Sutter et al., 1994
PLW-7	<i>p</i> <sub>syn</sub> (E/L)	MCS	Del III	Transient gpt <sup>e</sup>	Wyatt et al., 1996
PMC03	<i>p</i> <sub>syn</sub> (E/L)	MCS	Del III	GUS	Carroll and Moss, 1995
PLW-9	<i>pH5</i> (E/L)	MCS	Del III	Transient gpt <sup>e</sup>	Wyatt et al., 1996
PLW-17	<i>pH5</i> (E/L)	MCS	Del II	None	L. Wyatt and B. Moss, unpub. Observ.
PLW-21	<i>p</i> <sub>syn</sub> (E/L)	MCS	Del II	None	L. Wyatt and B. Moss, unpub. Observ.
PLW-22	<i>p</i> <sub>syn</sub> (E/L)	MCS	Del II	$\beta$ -gal	L. Wyatt and B. Moss, unpub. Observ.
PLW-24	<i>p7.5</i> (E/L)	MCS	Del II	None	L. Wyatt and B. Moss, unpub. Observ.

<sup>a</sup>pRB21 was specifically designed for use with vaccinia virus vRB12, which has a deletion in the F13L gene. The plasmids pG06, pLW-7, pMC03, pLW-9, pLW-17, pLW-21, pLW-22, and pLW-24 were designed for MVA.

<sup>b</sup>Abbreviations: E, early; L, late; E/L, early and late. The designation "x2" refers to two oppositely oriented promoters that can be used for expression of two genes.

<sup>c</sup>*Sma*I digestion gives a blunt end for cloning any fragment that has been blunt-ended. MCS signifies multiple cloning sites.

<sup>d</sup>Abbreviations: TK, thymidine kinase locus; F12L/F13L, between F12L and F13L open reading frames; Del III, site of natural deletion in MVA.

<sup>e</sup>Transient selection in which DXGPRT gene is deleted from recombinant vaccinia virus during recombination; see Background Information.

<sup>f</sup>Chakrabarti et al., 1997 *Biotechniques* 23:1094-1097; Sutter et al., 1994 *Vaccine* 12:1032-1040; Wyatt et al., 1996 *Vaccine* 14:1451-1458; Carroll and Moss 1995 *BioTechniques* 19:352-355.

[0031] The vaccinia virus expression vector of the present invention has utility for facilitating the expression of a broad number of exogenous nucleotide sequences. Genes, cDNAs encoding open reading frames derived from prokaryotic, eukaryotic, or viral sources other than vaccinia virus have been expressed using vaccinia virus vectors. The gene of interest is usually placed next to a vaccinia promoter and this expression cassette is then inserted into the virus genome by homologous recombination or direct ligation. Use of poxvirus promoters can be important since cellular and other viral promoters are not recognized by the vaccinia transcriptional apparatus. Strong late promoters are preferable when high levels of expression are desired. Early and intermediate-stage promoters, however, can also be used. The most versatile and widely used promoters contain early and late promoter elements. A list of suitable promoters appears in TABLE 1.

[0032] A number of plasmids are available with restriction endonuclease sites for insertion of foreign genes downstream of vaccinia promoters. The expression cassette is generally flanked by vaccinia DNA to permit homologous recombination when the plasmid is transfected into cells that have been infected with a vaccinia virus. The flanking vaccinia virus DNA is chosen so that recombination will not interrupt an essential viral gene. The shuttle vectors listed in TABLE 1 each have a multiple cloning site (MCS) to facilitate the insertion of exogenous nucleotide sequences.

[0033] In one embodiment, the vaccinia virus vectors of the present invention are contemplated for use as expression vectors *in vitro* and *in vivo*. Whole genes, open reading frames (ORFs), and other exogenous nucleotide fragments, such as nucleic acid sequences to generate antisense products, are contemplated for expression using the vaccinia virus vectors of the present invention.

[0034] Classes of genes contemplated for expression with the vectors of the present invention include tumor suppressor genes, cytotoxic genes, cytostatic genes, cytokines, and antigen encoding genes. Examples of tumor suppressor genes include: WT1, p53, p16, Rb, BRCA1, and others. Genes with therapeutic or scientific applications include: cystic fibrosis transmembrane regulator (CFTR), Factor VIII, low density lipoprotein

Station	Time	Lat	Long	Alt	Temp	Wind	Clouds	Remarks
1	0000	33° 15' N	118° 15' E	10	28.5	0	0	
2	0100	33° 15' N	118° 15' E	10	28.5	0	0	
3	0200	33° 15' N	118° 15' E	10	28.5	0	0	
4	0300	33° 15' N	118° 15' E	10	28.5	0	0	
5	0400	33° 15' N	118° 15' E	10	28.5	0	0	
6	0500	33° 15' N	118° 15' E	10	28.5	0	0	
7	0600	33° 15' N	118° 15' E	10	28.5	0	0	
8	0700	33° 15' N	118° 15' E	10	28.5	0	0	
9	0800	33° 15' N	118° 15' E	10	28.5	0	0	
10	0900	33° 15' N	118° 15' E	10	28.5	0	0	
11	1000	33° 15' N	118° 15' E	10	28.5	0	0	
12	1100	33° 15' N	118° 15' E	10	28.5	0	0	
13	1200	33° 15' N	118° 15' E	10	28.5	0	0	
14	1300	33° 15' N	118° 15' E	10	28.5	0	0	
15	1400	33° 15' N	118° 15' E	10	28.5	0	0	
16	1500	33° 15' N	118° 15' E	10	28.5	0	0	
17	1600	33° 15' N	118° 15' E	10	28.5	0	0	
18	1700	33° 15' N	118° 15' E	10	28.5	0	0	
19	1800	33° 15' N	118° 15' E	10	28.5	0	0	
20	1900	33° 15' N	118° 15' E	10	28.5	0	0	
21	2000	33° 15' N	118° 15' E	10	28.5	0	0	
22	2100	33° 15' N	118° 15' E	10	28.5	0	0	
23	2200	33° 15' N	118° 15' E	10	28.5	0	0	
24	2300	33° 15' N	118° 15' E	10	28.5	0	0	

**[0035]**

**[0036]**

**[0037]**

bystander effect is seen. Several investigators have studied this phenomenon because a strong bystander effect is thought to be essential *in vivo* when tumor cell transduction is inefficient. The *in vitro* bystander effect for the HSV-TK system is thought to be mediated by gap junctions which allow transport of GCV triphosphate between cells. The observation that cell lines with relatively few gap junctions had minimal bystander effects led to the upregulation of a gap junction protein (connexin) in a glioma cell line which increased the bystander effect *in vitro* and *in vivo*.

[0038] Several *in vivo* studies have demonstrated the potential therapeutic utility of this system. Subcutaneous sarcomas in BALB/c mice, retrovirally transduced with the HSV-TK gene, demonstrated a complete response after treatment with intraperitoneal (IP) GCV. Hepatic metastases transduced *in vivo* with a retrovirus expressing HSV-TK, were treated with twice daily injection of GCV for 5 days. This led to a significant inhibition of tumor growth compared to controls, with some complete regressions. Non-replicating, recombinant adenovirus carrying the HSV-TK gene has been used in many tumor models. Mice with intracerebral gliomas were injected intratumorally with adenovirus carrying HSV-TK, and then received IP GCV twice daily for 6 days. This resulted in a 23-fold decrease in tumor size compared to controls, and 2 complete regressions. There was no toxicity to the surrounding normal brain. Similar adenoviral/HSV-TK/GCV systems have effectively treated experimental ovarian cancer, head and neck cancer, breast cancer and melanoma. No systemic toxicity from the converted prodrug has been demonstrated.

[0039] The varicella thymidine kinase gene is another suicide gene contemplated for use with the present invention. Varicella zoster (VZV) is another virus in the *Herpesviridae* family, whose TK gene shares 28% homology with the HSV-TK and has a different substrate specificity. Several prodrugs have been tested for activity in a suicide gene system using VZV-TK, including GCV, acyclovir, (E)-5-(2-bromovinyl)-2'-deoxyuridine (BVDU), and others.

[0040] Breast cancer cells retrovirally transfected with VZV-TK, or controls, were treated with GCV, acyclovir, or BVDU. In the non-transduced controls there was minimal toxicity due to the prodrugs. In the transduced cell lines there was minimally increased sensitivity to GCV or acyclovir, but extreme sensitivity to BVDU (400 to

2000-fold over the parental cells). 50% of glial cells were needed to express the VZV-TK in order to see a global cytotoxic effect. The bystander effect was dependent on the cell line tested rather than the VZV-TK/BVDU system itself and may be a reflection of the number of intercellular gap junctions similar to the HSV-TK system.

[0041] Subcutaneously injected cells expressing the VZV-TK gene were similarly sensitive to BVDU. Significant, dose dependent, growth inhibition was seen after 8 daily IP injections of BVDU though no complete regressions were seen. There was no systemic toxicity from the converted prodrug.

[0042] Another suicide gene is cytosine deaminase (CD), which is a bacterial enzyme that deaminates cytosine to uracil and 5-fluorocytosine (5-FC) to 5-fluorouracil (5-FU). As it is not normally present in mammalian cells, the CD gene is useful in suicide gene systems. Transfer of the CD gene to mammalian tumor cells allows local conversion of 5-FC into 5-FU, which inhibits both RNA and DNA synthesis and leads to cell death. 5-FU is an effective chemotherapy agent and it has selectivity for dividing cells. It is standard therapy for various gastrointestinal and breast cancers. The prodrug 5-FC is well tolerated by humans and has been used in antifungal therapy.

[0043] Several methods to selectively transfer the CD gene to tumor cells have been employed including plasmid transfections, retroviral transduction, adenoviral infection, vaccinia infection and antibody-conjugates.

[0044] *In vitro* studies have demonstrated the ability to sensitize tumor cells to 5-FC by expression of the CD gene. The ability of 5-FU to diffuse through the cell membrane makes this an attractive system because of the bystander effect. When 20% of the cells expressed the CD gene, 60 to 80% cytotoxicity was seen after 4 days. The detection of 5-FU in the supernatants of CD-transduced cells suggests that direct cell to cell contact (such as in the TK systems) is not needed for a bystander effect.

[0045] Subcutaneous tumors, retrovirally transduced with the CD gene, showed significant growth inhibition which was further improved with earlier treatment and higher doses of 5-FC. The use of viruses to deliver the CD gene to subcutaneous xenografts has also been successful with significant growth inhibition or tumor regression when injected intra-tumorally. Intra-hepatic or intravenous delivery of an adenovirus carrying the CD gene



prolongation of survival with a 30% and 50% cure rate respectively. No evidence of toxicity from the converted prodrug was seen with this dosing regimen, but significant hepatic toxicity was seen at higher doses, presumably due to the potency and diffusability of 6-MP. This system is ultimately limited by systemic toxicity of the converted prodrug, requiring careful prodrug dosing.

$\beta$ -lactamase is a bacterial enzyme that confers resistance to  $\beta$ -lactam antibiotics by cleavage of the amide bond in the  $\beta$ -lactam ring. The synthesis of nontoxic prodrugs, consisting of cephalosporins conjugated to chemotherapy agents, has allowed the system to be used for suicide gene therapy as the conjugated prodrug is similarly cleaved releasing active drug. Several prodrugs have been synthesized for this system including cephalosporin-doxorubicin (C-Dox), 7-(4-carboxybutanamido)-cephalosporin mustard (CCM), cephalosporin-mitomycin C, cephalosporin-paclitaxel, and cephalosporin-carboplatinum. Tumor-specific antibodies conjugated to the  $\beta$ -lactamase enzyme have allowed for tumor-targeted activation of the various prodrugs.

[0049] *In vitro* studies showed that treatment of melanoma cells with an antibody- $\beta$ -lactamase conjugate followed by the prodrugs C-Dox and CCM, led to a 4 and 30-fold increase respectively, in the sensitivity of cells to the prodrugs although some nonspecific prodrug toxicity was seen. Subcutaneous melanoma xenografts in nude mice treated weekly with antibody- $\beta$ -lactamase and C-Dox showed tumor regression which eventually progressed. In contrast, similar treatments using CCM led to complete long-term regression in 80% of the mice. Although some neurologic toxicity was seen with the C-Dox, there was no systemic toxicity from the CCM. Similar studies using antibody- $\beta$ -lactamase/C-Dox to treat xenografts of breast carcinoma cells and colon carcinoma cells showed tumor growth inhibition in both histologies with minimal toxicity, although no complete regressions were seen. Finally, nude mice with subcutaneous melanoma xenografts were treated using a single-chain antibody/ $\beta$ -lactamase fusion protein in combination with CCM. Complete tumor regressions, in the majority of animals treated with CCM 12 hours after delivery of the fusion protein, were seen with no evidence of systemic toxicity. While this system has not been studied in a gene delivery approach, the  $\beta$ -lactamase protein is extremely versatile and should be functional in this context.

Accordingly, this gene may provide an excellent means by which to kill cells infected with the double mutant vaccinia virus of the present invention expressing this gene.

[0050] Carboxypeptidase G2 (CPG2) is a bacterial enzyme that catalyzes the hydrolytic cleavage of folates and methotrexate to pteroates and L-glutamic acid. Similar cleavage releases active alkylating agents from prodrugs synthesized with a glutamic acid group blocking the enzyme's function. Various methods of tumor specific expression of CPG2 have been studied including antibody targeting and stable transfection, suggesting it would be a good enzyme for use in a suicide gene system.

[0051] Stable transfection of colon and ovarian carcinoma cell lines with CPG2 resulted in a 16 to 95-fold increase in sensitivity to the prodrug 4-[(2-chloroethyl)(2-mesyloxyethyl) amino] benzoyl-L-glutamic acid (CMDA). 100% cytotoxicity was seen when 3.7 to 12% of the cells expressed CPG2. *In vivo* studies have used tumor specific monoclonal antibodies conjugated to CPG2 as a means of enzyme delivery. A nude mouse model of human choriocarcinoma showed significantly increased survival when treated with anti-human chorionic gonadotrophin (HCG) antibody/CPG2 conjugates, followed by only 3 doses of prodrug 72 hours later. Subcutaneous colon carcinoma xenografts had significant growth delay when treated with antibody/CPG2 conjugate followed by prodrug, although the tumors subsequently regrew. Minimal toxicity to the prodrug and its metabolites was seen.

[0052] Cytochrome P450-2B1 (CYP2B1) is a naturally occurring liver enzyme necessary in the conversion of cyclophosphamide and ifosfamide to their 4-hydroxy derivatives. These metabolites are unstable and further degrade to become the toxic metabolites acrolein and phosphoramidate mustard, which cause protein and DNA alkylation respectively. Because tumor levels of the enzyme are low, these chemotherapeutic agents are normally metabolized in the liver, leading to systemic toxicity prior to a maximal tumor response. Use in a suicide gene system allows upregulation of CYP2B1 expression in tumor cells, leading to local conversion of cyclophosphamide and ifosfamide, and potentially less systemic toxicity. This system is appealing because it may enhance the effect of agents which have already shown to be effective in cancer therapy and for which a safety profile has already been defined.



[0053] *In vitro* studies demonstrated that glioma cells and breast cancer cells, stably transfected with the CYP2B1 gene, acquired sensitivity to cyclophosphamide and ifosfamide. This sensitivity was blocked by the CYP2B1 inhibitor metyrapone. A significant bystander effect was seen when 20 - 50 % of the cells expressed the CYP2B1 gene, and this persisted in co-culture experiments suggesting that direct cell contact is not needed for this effect.

[0054] Several studies have shown enhanced sensitivity of subcutaneous tumors (glioma, breast cancer) retrovirally transduced with CYP2B1 (*ex vivo*) to cyclophosphamide. In one study, this led to complete inhibition of tumors in 95% of animals after a single IP injection of cyclophosphamide. Inoculation of retroviral producer cells or replication deficient adenovirus expressing the CYP2B1 gene into intracerebral gliomas led to prolonged survival after 1 dose of IP cyclophosphamide. There was no notable toxicity from the cyclophosphamide conversion at the doses studied.

[0055] Finally, Chase *et al* (1998) were able to show that intratumoral injection of a CYP2B1 expressing herpes simplex virus and cyclophosphamide, had an enhanced oncolytic effect compared to the virus alone. Chase M, et al., Nature Biotech 16:444-8 (1998).

[0056] Another interesting suicide gene is nitroreductase. Initial studies of the alkylating agent CB1954 showed promise in a rat Walker tumor model, however human trials were disappointing. Subsequently it was shown that the human equivalent of rat DT diaphorase, was inefficient at converting CB1954 to its active metabolite, the 4-hydroxylamino derivative which causes DNA crosslinking and cell death. The identification of a bacterial enzyme with similar activity, *E. coli* nitroreductase (ntr), led to its use in suicide gene systems, where tumor-specific expression of the bacterial enzyme would render tumor cells uniquely sensitive to CB1954.

[0057] *In vitro* studies have shown a 10 to 100-fold increase in sensitivity of pancreatic, colonic, and mammary cell tumor lines to CB1954 when transfected with plasmids expressing the ntr gene. Retroviral transduction led to a 17 to 500-fold increase insensitivity to CB1594, which correlated with the amount of nitroreductase produced by the cells. Cell mixing experiments demonstrated a significant *in vitro* bystander effect if 30% of

the cells express the gene. Transgenic mice engineered to express a CD2/ntr transgene in T cells and thymocytes had much smaller spleens and thymuses, and a significantly increased level of apoptosis in these organs, 5 days after treatment with CB1954. Clark *et al* (1997) showed that transgenic mice expressing ntr under the control of the  $\beta$ -lactoglobulin promoter, had increased levels of ntr mRNA and protein in mammary glands compared to other tissues, and treatment with CB1954 resulted in disrupted mammary glands with increased apoptosis. Clark et al., Gene Ther. 4:101-10 (1997). Nonspecific toxicity from the prodrug or converted active drug was not seen.

[0058] Traditionally the *E. coli* xanthine-guanine phosphoribosyl transferase (gpt) gene has been used for positive selection in a purine salvage pathway as it catalyzes the conversion of xanthine, hypoxanthine and guanine to their respective monophosphates. Cells which express the gpt gene can utilize xanthine as a purine analog in the presence of mycophenolic acid and hypoxanthine which inhibit the mammalian pathway. Conversely, gpt has recently been used to selectively sensitize tumor cells to 6 - thioxanthine (6-TX) as it is incorporated into this purine synthesis pathway. Mroz and Moolton showed that retrovirally transducing the gpt gene into sarcoma cells conferred an 86-fold increased sensitivity to 6-TX. . Human Gene Therapy 1993; 4:589-95. Tamiya *et al* showed that glioma cells were sensitive to 6-TX at an LD50 of 2.5  $\mu$ mol/L when transduced with gpt. Cancer Gene Therapy 1996; 3:155-62 Untransfected cells were minimally sensitive at 50  $\mu$ mol/L. As well, a significant bystander effect (75% cytotoxicity) was seen when 10% of the cells expressed gpt. This effect was abrogated when the cells were separated suggesting that cell to cell contact was required. Syngeneic Balb/C mice injected subcutaneously with a transduced sarcoma cell line showed complete regression in 95% of the animals after 5 days of treatment with 6-TX. A 10% treatment mortality due to toxicity of the converted prodrug was seen and was eradicated with an alternate day dosing schedule. Nude mice subcutaneously injected with transduced glioma cells showed a significant inhibition of tumor growth when treated with IP 6-TX for 10 days at a reduced dose, with no treatment toxicity. In the same study, nude mice with transduced intracerebral gliomas showed prolonged survival after 6-TX treatment. In another study similar subcutaneous tumor inhibition was seen, and regrowth was correlated with loss of the retrovirally expressed gpt

[illegible][illegible][illegible][illegible]

This leads to tumor specific toxicity when delivered via antibody-targeting or retroviral transduction, with no noted systemic toxicity.

[0061] Bovine carboxypeptidase A has been shown to cleave the amide bond of a methotrexate-phenylalanine prodrug, yielding free methotrexate. This enzyme was conjugated to a monoclonal antibody for tumor specific delivery, and *in vitro* studies showed that ovarian carcinoma cells became sensitive to the methotrexate-phenylalanine prodrug in the presence of carboxypeptidase A. Finally, a novel rabbit cytochrome P450 isoenzyme (CYP4B1) has been shown to convert the prodrugs 2-aminoanthracene (2-AA) and 4-ipomeanol (4-IM) into toxic alkylating agents. Glioma cells stably transfected with CYP4B1 showed increased sensitivity to 2-AA and 4-IM *in vitro* and *in vivo* compared to control. A strong bystander effect was seen with 70 to 80% cytotoxicity when 1% of the cells expressed the CYP4B1 gene.

[0062] A table of certain suicide genes and drugs with activity associated with these genes is found at TABLE 2.

TABLE 2

Enzyme/Prodrug Systems

ENZYME	PRODRUG	ACTIVE DRUG
<i>Herpes Simplex Virus</i> thymidine kinase	Gancyclovir	Gancyclovir triphosphate
<i>Varicella Zoster Virus</i> thymidine kinase	(E)-5-(2-bromovinyl)-2'-deoxyuridine (BVDU)	BVDU triphosphate
Cytosine deaminase	5-fluorocytosine	5-fluorouracil
Purine nucleoside phosphorylase	6-methylpurine deoxyriboside	6-methylpurine
$\beta$ -lactamase	7-(4-carboxybutanamido)-cephalosporin mustard	phenylenediamine mustard
Carboxypeptidase G2	4-[(2-chloroethyl)(2-mesyloxyethyl)amino] benzoyl-L-glutamic acid (CMDA)	4-[(2-chloroethyl)(2-mesyloxyethyl)amino] benzoic acid (CJS11)
Cytochrome P450-2B1	Cyclophosphamide/ifosfamide	acrolein + phosphoramidate mustard
<i>E. coli</i> nitroreductase	CB1954 (S-aziridin-yl-2-4-dinitrobenzamide)	5-aziridin-1-yl-4-hydroxylamino-2-nitrobenzamide
Xanthine-guanine phosphoribosyl-transferase	6-thioxanthine	6-thioxanthine monophosphate
$\beta$ -glucuronidase	epirubicin-glucoronide	Epirubicin

ENZYME	PRODRUG	ACTIVE DRUG
Thymidine phosphorylase	5'-deoxy-5-fluorouridine	5-fluorouracil
Deoxycytidine kinase	Cytosine arabinoside	Cytosine ababinoside monophosphate
Carboxylesterase	7-ethyl-10-[4-(1-piperidino)-1-piperidino] carbonyloxycamptothecin (CPT-11)	7-theyl-10- hydroxycamptothecin (SN-38)
Linamarase/ $\beta$ -glucosidase	Linamarin/Amygdalin	Cyanide
Carboxypeptidase A	Methotrexate-phenylalamine	Methotrexate
Cytochrome P450-4B1	2-aminoanthracene, ipomeanol	4- unknown alkylating agents

[0063] The vaccinia virus expression vectors of the present invention are also contemplated for *in vitro* expression purposes. Any gene, ORF, or other segment of nucleic acid can be expressed *in vitro* using the expression vectors of present invention.

[0064] The present invention has utility in expressing genes in vivo and in vitro. Examples of genes expressable by the vaccinia virus expression vector of the present invention include expressing human genes. An exemplary list of genes includes the list of human genes and genetic disorders authored and edited by Dr. Victor A. McKusick and his colleagues at Johns Hopkins and elsewhere, and developed for the World Wide Web by NCBI, the National Center for Biotechnology Information. Online Mendelian Inheritance in Man, OMIM (TM). Center for Medical Genetics, Johns Hopkins University (Baltimore, MD) and National Center for Biotechnology Information, National Library of Medicine (Bethesda, MD), 1999. << <http://www.ncbi.nlm.nih.gov/omim/>>>. These genes include: 239f2h9, 3pk, 4ebp1, 4ebp2, a11, a12m1, a12m2, a12m3, a12m4, a15, a1b, a1bg, a1st, a2m, a2mr, a2mrap, aa, aaa, aaa, aabt, aac1, aac2, aact, aadac, aanat, aars, aas, aat, aavs1, abc1, abc2, abc3, abc7, abc8, abcr, abi1, abl1, abl2, abll, abo, abp, abp1, abpa, abpx, abr, acaa, acac, acaca, acacb, acadl, acadm, acads, acadsb, acadvl, acat, acat1, acat2, acc, accb, accn1, accn2, accpn, ace1, ach, ache, achm1, achm2, achrb, achrd, achrg, acis, acly, aco1, aco2, acox, acox1, acox2, acox3, acp1, acp2, acp5, acpp, acr, acrv1, acs3, acs3, acs4, act2, act35, acta1, acta2, acta3, actb, actc, actg1, actg2, actl1, actn1, actn2, actn3, actsa, acug, acvr1, acvr2b, acvr11, acvrk1, acvrk2, acvrk3, acy1, ad1, ad2, ad3, ad4, ad5, ada, adam10, adam11, adam12, adam3, adam3a, adam3b, adam8, adar, adarb1, adarb2, adcp1, adcp2, adcy1, adcy2, adcy3, adcy3, adcy4, adcy5, adcy6, adcy7, adcy8, adcy9, adcyap1, adcyap1r1, add1, add2, add3, addl, adfn,

adh1, adh2, adh3, adh4, adh5, adh7, adhaps, adhcl@, adhr, adhr, adk, adl, adm, admix, adora1, adora2a, adora2b, adora2l, adora2l1, adora3, adprt, adra1a, adra1b, adra1c, adra1d, adra2a, adra2b, adra2c, adra2l1, adra2l2, adra2r, adrb1, adrb1r, adrb2, adrb2r11, adrb3, adrbk1, adrbk2, adsl, adss, adtb1, adx, adxr, ae1, ae2, ae3, aegl1, aemk, aes, afl0, afl7, af4, af6, af8t, af9, afd1, afdn, afg3, afg3l1, afm, afp, afx1, aga, agc1, ager, agl, agmx1, agmx2, agp1, agp7, agps, agrn, agrp, agrt, ags, agt, agtil, agr1, agr1a, agr2, agr11, agxt, ahc, ahcy, ahd, ahds, ahnak, aho2, ahr, ahsg, ahx, aib1, aic, aicl, aied, aih1, aih2, aih3, aim1, air, airc, aire, ak1, ak2, ak3, akap149, akt1, akt2, aku, alad, alas1, alas2, alb, alb2, alba, alcam, ald, aldh1, aldh10, aldh2, aldh3, aldh4, aldh5, aldh6, aldh9, ald11, aldoa, aldob, aldod, aldr1, alds, alk, alk1, alk2, alk3, alk6, alms1, alox12, alox15, alox5, alp, alpi, alpl, alpp, alpp12, alr, alr, als1, als2, als4, als5, alss, ambn, ambp, amcd1, amcd2b, amcn, amcn1, amcx1, amd1, amdm, amelx, amely, amfr, amg, amgl, amgx, amh, amhr, amhr2, aml1, aml1t1, aml2, aml3, amog, ampd1, ampd2, ampd3, amph, amph1, ampk, amt, amyla, amy1b, amy1c, amy2a, amy2b, an2, anc, ancr, ang, ang1, anh1, ank1, ank2, ank3, anop1, anova, anp, anpep, anpra, anprb, anprc, ans, ant1, ant2, ant3, ant3y, anx1, anx11, anx13, anx2, anx214, anx3, anx4, anx5, anx6, anx7, anx8, aoah, aoc2, aox1, ap2tf, apah1, apba1, apba2, apbb1, apbb2, apc, apcs, ape, apeced, apeh, apex, api1, api2, api3, apj, aplp, aplp1, aplp2, apnh, apo3l, apoa1, apoa2, apoa4, apob, apobec1, apoc1, apoc2, apoc3, apoc4, apod, apoe, apoer2, apoh, apolmt, apolp1@, apolp2@, app, appbp1, appl1, aprf, aprt, aps, apt1, apt1lg1, apx1, apy, aqdq, aqp0, aqp1, aqp2, aqp2l, aqp3, aqp4, aqp5, aqp6, aqp7, ar, ar1, ara, araf1, araf2, arcn1, ard1, ard1, areg, arf1, arf2, arf3, arf4l, arf5, arg, arg1, args, arh12, arh6, arh9, arha, arhb, arhc, arhg, arhgap2, arhgap3, arhgap6, arhgdia, arhgdib, arhh, arix, arl2, armd1, arnt, arntl, aro, arp, arp1, arpkd, arr3, arrb1, arrb2, arsa, arsacs, arsb, arsc1, arsc2, arsd, arse, arsf, art, art1, art3, art4, arts, arvd1, arvd2, arvd3, arvd4, as, asat, asb, ascl1, ascl2, asct1, asd1, asd2, asgr1, asgr2, ash1, asip, asl, asln, asm1, asma, asmd, asmt, asmtlx, asmtly, asnrs, asns, aspa, ass, astml, astn, asv, at, at1, at2r1, at3, ata, atbf1, atcay, atf1, ath1, aths, atm, atoh1, atox1, atp1a1, atp1a2, atp1a3, atp1a11, atp1b1, atp1b2, atp1b3, atp1b11, atp1g1, atp2a1, atp2a2, atp2a3, atp2b, atp2b1, atp2b2, atp2b2, atp2b3, atp2b4, atp4a, atp4b, atp5, atp5a, atp5b, atp5g1, atp5g2, atp5g3, atp5o, atp6a, atp6b1, atp6c, atp6e, atp6n1, atp7a, atp7b, atpm, atpsb, atpsk1, atpsk2, atq1, atr, atr, atr1, atr1, atr2, atrc1, atrc2, atrx, ats, atsv, atx1, atx2, au, aufl, aufla,

aut, avcd, aved, avp, avpr1a, avpr1b, avpr2, avpr3, avrp, avsd, awal, axl, axllg, axsf, azf1, azf2, azgp1, azul, b120, b144, b1g1, b29, b2m, b2mr, b3galt4, b4galt1, ba2r, babl, bag1, bai1, bai2, bai3, bak1, bam22, bap1, bap135, bapx1, bard1, bark2, bas, bat1, bat2, bat3, bat4, bat5, bax, bb1, bbbg1, bbbg2, bbs1, bbs2, bbs3, bbs4, bbs5, bcas1, bcat1, bcat2, bcate2, bcd1, bcei, bche, bckdha, bckdha, bcl1, bcl10, bcl2, bcl2a1, bcl2l2, bcl3, bcl5, bcl6, bcl7, bcl7a, bcl8, bcl9, bclw, bcm, bcm1, bcma, bcns, bcns, bcp, bcpm, bcpr, bcr, bcr12, bcr13, bcr14, bcsg1, bct1, bct2, bdb, bdb1, bdc, bde, bdkrb1, bdkrb2, bdmf, bdmr, bdnf, bed, bedp, bek, bene, bevi, bf, bf1, bf2, bfhd, bfic, bfls, bfnc2, bfp, bfsp1, bft, bglap, bgmr, bgn, bgp, bhd, bhpcdh, bhr1, bcd1, bid, bigh3, bin1, bir, bjs, bkma1, blast1, blau, blk, blm, blmh, bltr, blvra, blvrb, blym, bmal1, bmd, bmh, bml1, bmp1, bmp2, bmp2a, bmp2b1, bmp3, bmp4, bmp5, bmp6, bmp7, bmp8, bmp1a, bmp1b, bmx, bmyb, bn51t, bnc, bnc1, bnp, bor, bpad, bpag1, bpag2, bpes, bpes1, bpes2, bpgm, bphl, bpi, br, br140, braf, brca1, brca2, brca3, brcacox, brcd1, brcd2, brdt, brf1, brhc, bric, brks, brn3a, brn3b, brn3c, brn1, brw1c, bs, bsap, bsep, bsf2, bsg, bsnd, bssl, bst1, bst2, btak, btc, btd, bteb, bteb1, btg1, btg2, bths, btk, btkl, btn, bts, bub1b, bubr1, bwrla, bwrlb, bws, bwscr1a, bwscr1b, bzip, bzx, c11orf13, c1nh, c1qa, c1qb, c1qbp, c1qg, c1r, c1s, c2, c21orf1, c21orf2, c21orf3, c2ta, c3, c3br, c3dr, c3g, c4a, c4b, c4bpa, c4bpb, c4f, c4s, c5, c5ar, c5r1, c6, c7, c8a, c8b, c8g, c9, ca1, ca12, ca125, ca2, ca21h, ca3, ca4, ca5, ca6, ca7, ca8, ca9, caaf1, cabp9k, cac, cac@, caca, cacd, cacna1a, cacna1b, cacna1c, cacna1d, cacna1e, cacna1f, cacna1s, cacna2, cacnb1, cacnb2, cacnb3, cacnb4, cacng, cacn11a1, cacn11a2, cacn11a3, cacn11a4, cacn11a5, cacn11a6, cacn12a, cacn1b1, cacnlg, cact, cacy, cad, cad11, cadasil, cae1, cae3, caf, cafla, caga, cagb, cain, cak, cak1, cal11, calb1, calb2, calb3, calc1, calc2, calca, calcb, calcr, cald1, calla, calm1, calm2, calm3, calm11, calm13, calna, calna3, calnb, calnb1, calr, cals, calt, calu, cam, camk4, camkg, caml1, camlg, camp, can, canp3, canx, cap2, cap3, cap37, capb, capg, capl, capn1, capn2, capn3, capn4, cappa2, cappb, capr, caps, capza2, capzb, car, carp, cars, cart1, cas, cas2, casil, casp1, casp10, casp2, casp3, casp3, casp4, casp5, casp6, casp7, casp8, casq1, casq2, casr, cast, cat, cat1, cat4, catf1, catm, cav1, cav2, cav3, cbbm, cbd, cbfa1, cbfa2, cbfa2t1, cbfa3, cbfb, cbg, cbl, cbl2, cbln2, cbp, cbp, cbp2, cbp68, cbr1, cbs, cbt, cbt1, cc10, cca, cca1, cca11, cca12, ccb11, ccckr5, ccg1, ccg2, cchl1a1, cchl1a2, cchl1a3, cchl1b1, cck, cckar, cckbr, ccl, ccm1, ccm2, ccm3, ccn1, ccna, ccnb1, ccnc, ccnd1, ccnd2, ccnd3, ccne, ccnf, ccng1, ccnh,





cnn3, cnp, cnr1, cnsn, cntf, cntfr, cntn1, co, cocal, coca2, coch, cod1, cod2, coh1, coil, coll0a1, coll1a1, coll1a2, coll2a1, coll3a1, coll5a1, coll6a1, coll7a1, coll8a1, coll9a1, colla1, colla2, collar, col2a1, col3a1, col4a1, col4a2, col4a3, col4a4, col4a5, col4a6, col5a1, col5a2, col6a1, col6a2, col6a3, col7a1, col8a1, col8a2, col9a1, col9a1, col9a2, col9a3, colq, comp, comt, copeb, copt1, copt2, cord1, cord2, cord5, cord6, cort, cot, cox10, cox4, cox5b, cox6a1, cox6b, cox7a1, cox7a2, cox7a3, cox7am, cox8, cp, cp107, cp115, cp20, cp47, cp49, cpa1, cpa3, cpb2, cpb2, cpd, cpe, cpetr2, cpm, cpn, cpn1, cpn2, cpo, cpp, cpp32, cpp32, cppi, cps1, cpsb, cpsd, cpt1a, cpt1b, cpt2, cpu, cpx, cpx, cpxd, cr1, cr2, cr3a, crabp1, crabp2, crapb, crarf, crat, crbp1, crbp2, crd, crd1, creb1, creb2, crebbp, creb1, crem, crfb4, crfr2, crh, crhbp, crhr, crhr1, crhr2, crip, crk, crkl, crml, crmp1, crmp2, crp, crp1, crs, crs1c, crs2, crs3, crsa, crt, crtl1, crtm, crx, cry1, cry2, crya1, crya2, cryaa, cryab, cryb1, cryb2, cryb3, cryba1, cryba2, cryba4, crybb1, crybb2, crybb3, cryg1, cryg2, cryg3, cryg4, cryg8, cryg@, cryga, crygb, crygc, crygd, crygs, crym, cryz, cs, csa, csb, csbp1, csci, csd, csd2, csda, cse, cse1, csf1, csf1r, csf2, csf2ra, csf2rb, csf2ry, csf3, csf3r, csh1, csh2, csk, csmf, csn1, csn10, csn2, csn3, csnb1, csnb2, csnb3, csnk1a1, csnk1d, csnk1e, csnk1g2, csnk2a1, csnk2a2, csnk2b, csnu3, cso, cspb, cspg1, cspg2, cspg3, csr, csrb, csrp, csrp1, csrp2, cst1, cst1, cst2, cst3, cst4, cst5, cst6, csta, cstab, csx, ct2, ctaa1, ctaa2, ctag, ctb, ctbp1, ctbp2, ctgf, cth, cthm, ctk, ctla1, ctla3, ctla4, ctla8, ctm, cttna1, cttna2, cttnb1, cttnnd, cttnnd1, cttnr, cttns, ctp, ctpct, ctps, ctrl, ctr2, ctrb1, ctrl, ctsa, ctsb, ctsc, ctsd, ctse, ctsg, ctsgl2, ctsh, ctsk, ctsl, ctss, ctsw, ctsz, ctx, cubn, cul3, cul4b, cul5, cutl1, cvap, cvd1, cvl, cx26, cx31, cx32, cx37, cx40, cx43, cx46, cx50, cxb3s, cxcr4, cxorf4, cyb5, cyb561, cyba, cybb, cycl, cyk4, cyld1, cyp1, cyp11a, cyp11b1, cyp11b2, cyp17, cyp19, cypl1a, cypl1a2, cypl1b1, cyp21, cyp24, cyp27, cyp27a1, cyp27b1, cyp2a, cyp2a3, cyp2a6, cyp2b, cyp2c, cyp2c19, cyp2c9, cyp2d, cyp2d@, cyp2e, cyp2e1, cyp2f1, cyp2j2, cyp3a4, cyp4a1, cyp4b1, cyp51, cyp7, cyp7a1, cyr61, cym1, cym2, czp3, d10s105e, d10s170, d10s170, d11s302e, d11s636, d11s813e, d11s833e, d12s2489e, d12s53e, d13s1056e, d13s25, d14s46e, d15s12, d15s226e, d15s227e, d16s2531e, d16s469e, d17s136e, d17s811e, d18s892e, d19s204, d19s381e, d1s111, d1s155e, d1s166e, d1s1733e, d1s2223e, d1s61, d2h, d2s201e, d2s448, d2s448e, d2s69e, d3s1231e, d3s1319e, d3s48e, d4, d4s90, d5s1708, d5s346, d6, d6s1101, d6s207e, d6s2244e, d6s2245e, d6s228e, d6s229e, d6s230e, d6s231e, d6s51e, d6s52e, d6s54e, d6s81e, d6s82e, d7s437, d8s2298e,





fpsl3, fpsl4, fpsl5, fr, frap1, fraxa, fraxe, fraxf, frda, freac2, freac6, freac9, frgl, frp1, frv1, frv2, frv3, fsg1, fsgs, fshb, fshd1a, fshmd1a, fshprh1, fshr, fssv, fth1, fthl6, ftl, ftz1, ftzfl, fuca1, fuca2, fur, fus, fuse, fut1, fut2, fut3, fut4, fut5, fut6, fut7, fut8, fvt1, fxr1, fxy, fy, fyn, fzd1, fzd2, fzd3, fzd5, fzd6, fzd7, fzf, g0s8, g10p1, g10p2, g17, g17p1, g19p1, glp1, glp2, glp3, g22p1, g6pc, g6pd, g6pd1, g6pdl, g6pt, g6pt1, g6s, g7p1, ga2, gaa, gabatr, gabpa, gabpb1, gabra1, gabra2, gabra3, gabra4, gabra5, gabra6, gabrb1, gabrb2, gabrb3, gabrd, gabre, gabrg1, gabrg2, gabrg3, gabrr1, gabrr2, gad1, gad2, gad3, gadd153, gadd45, gak, gal, galbp, galc, gale, galgt, galk1, galk2, galn, galnact, galnr, galnr1, galns, galnt1, galnt2, galnt3, galr1, galt, gan, gan1, ganab, ganc, gap, gap1m, gap43, gapd, gar22, garp, gars, gart, gas, gas1, gas2, gas41, gas6, gas7, gasr, gast, gata1, gata2, gata3, gata4, gata6, gay1, gba, gbas, gbbb1, gbbb2, gbel, gbp1, gbx2, gc, gcap, gcap2, gcdh, gcfl, gcf2, gcfx, gcg, gcgr, gch1, gck, gckr, gcn5l1, gcn5l2, gcnf, gcnt1, gcnt2, gcp, gcp2, gcs, gcs1, gcsf, gcsfr, gcsp, gctg, gcy, gda, gde, gdf5, gdf8, gdh, gdi1, gdi2, gdid4, gdld, gdnf, gdnfr, gdnfra, gdnfrb, gdx, gdxxy, ge, gem, geney, gey, gfl, gfl, gfap, gfer, gfer, gfi1, gfpt, gfra1, gfra2, ggcx, ggt1, ggt2, ggta1, ggta1, ggta2, gh1, gh2, ghc@, ghdx, ghn, ghr, ghfr, ghrh, ghrhr, ghs, ghv, gif, gifb, gip, gip, gipr, girk1, girk2, girk3, girk4, gja1, gja3, gja4, gja5, gja8, gjb1, gjb2, gjb3, gk, gk2, gla, glat, glb1, glb2, glc1a, glc1b, glc1c, glc1d, glc1f, glc3a, glc3b, glc1c, glc1r, glct2, glct3, glde, glepp1, glg1, gli, gli2, gli3, gli4, glnn, glns, glo1, glo2, glp1r, glra1, glra2, glra3, glrb, glrx, gls, glud1, glud2, glul, glur1, glur2, glur3, glur4, glur5, glur6, glur7, glut1, glut2, glut3, glut4, glut5, glvr1, glvr2, gly96, glya, glyb, glys1, glyt1, glyt1, glyt2, gm2a, gma, gmcsf, gmds, gml, gmpr, gmps, gna11, gna15, gna16, gnai1, gnai2, gnai2a, gnai2b, gnai2l, gnai3, gnai, gnaol, gnaq, gnas, gnas1, gnat1, gnat2, gnaz, gnb1, gnb2, gnb3, gng5, gnl1, gnpta, gnrh1, gnrh2, gnrhr, gns, gnt1, golga4, got1, got2, gp130, gp1ba, gp1bb, gp2, gp2b, gp39, gp3a, gp75, gp78, gp9, gpa, gpam, gpat, gpb, gpc, gpc1, gpc3, gpc4, gpd, gpd1, gpd2, gpds1, gpe, gpi, gpi2, gpm6a, gpm6b, gpoa, gpr1, gpr10, gpr11, gpr12, gpr13, gpr15, gpr17, gpr18, gpr19, gpr2, gpr20, gpr21, gpr22, gpr23, gpr25, gpr29, gpr3, gpr30, gpr31, gpr32, gpr35, gpr37, gpr39, gpr4, gpr5, gpr6, gpr7, gpr8, gpr9, gprcy4, gprk2l, gprk4, gprk5, gprk6, gprv28, gpsa, gpse, gpt, gpx1, gpx2, gpx3, gpx4, gr2, grb1, grb10, grb2, grf2, gria1, gria2, gria3, gria4, grid2, grik1, grik2, grik3, grik4, grik5, grin1, grin2a, grin2b, grin2c, grin2d, grina, grk1, grk5, grk6, grl, grll1, grm3, grm8, grmp, grn, gro1, gro2, gro3, grp, grp58, grp78,

grpr, grx, gs, gsl, gsas, gsc, gscl, gse, gshs, gsl, gsm1, gsn, gsp, gspt1, gsr, gss, gst12, gst11, gst2, gst2, gst3, gst4, gst5, gsta1, gsta2, gstm1, gstm11, gstm2, gstm3, gstm4, gstm5, gstpl1, gstt2, gt1, gt335, gta, gtb, gtbp, gtd, gtf2e2, gtf2f1, gtf2h1, gtf2h2, gtf2h4, gtf2i, gtf2s, gtf3a, gtg, guc1a2, guc1a3, guc1b3, guc2c, guc2d, guc2f, guca1a, guca1b, guca2, guca2, guca2a, guca2b, gucsa3, gucsb3, gucyl1a2, gucyl1a3, gucyl1b3, gucyl2c, gucyl2d, gucyl2f, guk1, guk2, gulo, gulop, gusb, gusm, gust, gxp1, gypa, gypb, gypc, gype, gys, gys1, gys2, gзма, gзмb, gзмh, gзмm, h, h142t, h19, h1f0, h1f1, h1f2, h1f3, h1f4, h1f5, h1fv, h2a, h2ax, h2az, h2b, h2b, h3f2, h3f3b, h3ft, h3t, h4, h4f2, h4f5, h4fa, h4fb, h4fc, h4fd, h4fe, h4fg, h4fh, h4fi, h4fj, h4fk, h4fl, h4fm, h4m, h6, ha2, habp1, hadha, hadhb, hadhsc, haf, hagh, hah1, haip1, hal, hap, hap1, hap2, hars, has2, hat1, hausp, hb1, hb1, hb6, hba1, hba2, hbac@, hbb, hbbc@, hbd, hbe1, hbegf, hbf2, hbg1, hbg2, hbgr, hbhr, hbm, hbp, hbq1, hbz, hc2, hc3, hca, hcat2, hccs, hcdh, hcf2, hcfcl, hcg, hck, hcl1, hcl2, hcl3, hclsl, hcp, hcp1, hcs, hcvs, hd, hdac1, hdc, hdgf, hdhc7, hdlbp, hdld, hdldt1, hdr, hed, hed, hegfl, hek, hek3, heln1, hem1, hema, hemb, hemc, hempas, hen1, hen2, hep, hep10, her2, her4, herg, herv1, hes1, hesx1, het, hexa, hexb, hf1, hf10, hfc1, hfe, hfe2, hfh11, hfsp, hgd, hgf, hgf, hgfl, hgl, hh, hh72, hhcl, hhcl2, hhd, hhh, hhmjg, hhr23a, hht1, hht2, hiap2, higm1, hilda, hint, hiomt, hip, hip1, hip116, hip2, hir, hira, his1, his2, hive1, hivep1, hivep2, hjcd, hk1, hk2, hk3, hk33, hke4, hke6, hkr1, hkr2, hkr3, hkr4, hl11, hl19, hla-a, hla-b, hla-c, hla-cda12, hla-dma, hla-dmb, hla-dna, hla-dob, hla-dpal, hla-dpb1, hla-dqa1, hla-dr1b, hla-dra, hla-e, hla-f, hla-g, hla-ha2, hladp, hlaf, hlals, hlcs, hlm2, hlp, hlp3, hlr1, hlr2, hlt, hlxl, hlxb9, hmaa, hmab, hmat1, hmbs, hmcs, hmg1, hmg14, hmg17, hmg2, hmgcl, hmgcr, hmgcs1, hmgcs2, hmgic, hmgiy, hmgx, hmmr, hmn2, hmox1, hmox2, hmr, hms1, hmsnl, hmx1, hmx2, hnd, hnfla, hnf2, hnf3a, hnf3b, hnf4a, hnp36, hnpcc6, hnrpa1, hnrpa2b1, hnrpd, hnrpf, hnrpg, hnrph1, hnrph2, hnrph3, hnrpk, homg, hops, hox10, hox11, hox12, hox1@, hox1a, hox1b, hox1c, hox1d, hox1e, hox1f, hox1g, hox1h, hox1i, hox1j, hox2@, hox2a, hox2b, hox2c, hox2d, hox2e, hox2f, hox2g, hox2h, hox2i, hox3@, hox3a, hox3b, hox3c, hox3d, hox3e, hox3f, hox3g, hox4@, hox4a, hox4b, hox4c, hox4d, hox4e, hox4f, hox4g, hox4h, hox4i, hox7, hox8, hoxa1, hoxa10, hoxa11, hoxa13, hoxa3, hoxa4, hoxa5, hoxa6, hoxa7, hoxa9, hoxa@, hoxb1, hoxb2, hoxb3, hoxb4, hoxb5, hoxb6, hoxb7, hoxb8, hoxb9, hoxb@, hoxc12, hoxc13, hoxc4, hoxc5, hoxc6, hoxc8, hoxc9, hoxc@, hoxd1, hoxd10, hoxd11, hoxd12, hoxd13, hoxd3, hoxd4, hoxd8,





mac2, mac25, macam1, macs, mad, mad211, madd, madh1, madh2, madh3, madh4, madh5, madh6, madh6, madh7, madh9, madm, madr1, maf, mafd1, mafd2, mag, mage1, mageb3, mageb4, magell, magoh, magp, magp1, magp2, mak, mal, mall, man2a2, manal, mana2, mana2x, manb, manb1, manba, maoa, maob, map1a, map1alc3, map1b, map1blc3, map2, map4, map80, map97, mapk1, mapkap3, mapkkk4, mapt, mar, mark3, mars, mas1, masp1, mat1a, mat2a, mata1, mata2, matk, matn1, matn3, max, maz, mb, mbd1, mbl, mbl2, mbp, mbp1, mbs, mbs2, mc1r, mc2r, mc3r, mc4r, mc5r, mcad, mcc, mcdcl, mcdrl, mcf2, mcf3, mcfdl, mch2, mch3, mch4, mch5, mckd, mcl, mcl1, mcm, mcm2, mcm2, mcm3, mcm6, mcm7, mcmt, mcop, mcor, mcp, mcp1, mcp3, mcph1, mcr, mcs, mcsf, mcspl, mct1, md1, mdb, mdc, mdcrl, mddc, mdeg, mdf1, mdg, mdg1, mdh1, mdh2, mdk, mdk, mdm2, mdm4, mdr1, mdr3, mdrsl, mdrv, mds, mds1, mdul, mdu2, mdu3, mdx, me1, me2, mea, mea6, mecl1, mecp2, med, mef, mef2a, mef2b, mef2c, mef2d, mefv, mehmo, meis1, meis2, mekk, mekk1, mekk4, mel, mel18, melf, memo1, men1, men2a, meox1, meox2, mep1a, mep1b, mer2, mer6, mest, met, metrs, mfap1, mfap2, mfap3, mfap4, mfd1, mfi2, mfs1, mfs2, mft, mfts, mg50, mga, mga1, mga3, mgat1, mgat2, mgat5, mgcl, mgcn, mgcr, mgct, mgdf, mgea, mgf, mgi, mgmt, mgp, mgsa, mgst1, mgst2, mhc, mhc2ta, mhp2, mhs, mhs2, mhs3, mhs4, mhs6, mia, mic10, mic11, mic12, mic17, mic18, mic2, mic2x, mic2y, mic3, mic4, mic7, mica, micb, mid1, midas, mif, mif, mig, mip, mip2a, mip2b, mip3b, mipep, mitf, miwc, mjd, mk, mki67, mkks, mkp2, mkp3, mkpx, mks, mks, mks1, mks2, mla1, mlck, mlf1, mlf2, mlh1, mlk1, mlk3, mll, mll2, mllt1, mllt2, mllt3, mllt4, mllt6, mllt7, mlm, mlm, mln, mlp, mlr, mlrg, mlrw, mls, mltn, mlvar, mlvi2, mlvt, mmac1, mme, mmp1, mmp10, mmp11, mmp12, mmp13, mmp14, mmp15, mmp16, mmp17, mmp19, mmp2, mmp21, mmp22, mmp3, mmp7, mmp8, mmp9, mn, mn, mnb, mnbh, mnda, mnng1, mnk, mns, mnt, mocod, mocs1, mocs2, mody1, mody3, mog, mok2, mom1, mos, mot2, mov34, mox1, mox2, mox44, moz, mp19, mpb1, mpd1, mpdz, mpe, mpe16, mpg, mpi, mpif2, mpl, mpllg, mpo, mpp1, mpp2, mpp3, mppb, mpri, mprm, mps2, mps3a, mps3c, mps4a, mpsh, mpts, mpv17, mpz, mr1, mr77, mrbc, mrc1, mre11, mre11a, mrg1, mrgh, mros, mrp, mrp, mrp1, mrpl23, mrs, mrsl, mrsr, mrst, mrx1, mrx14, mrx2, mrx20, mrx21, mrx23, mrx29, mrx41, mrx48, mrx49, mrx9, mrxa, mrxs1, mrxs2, mrxs3, mrxs4, mrxs5, mrxs6, mrxs8, ms3315, ms336, msg1, msh2, msh3, msh4, msh6, msil, msk16, msk39, msk41, mslr1, msmb, msn, msr1, mssl,



mss4, mss4, msse, mst, mstl, mstlr, mstd, mstn, msud1, mxsl, mxs2, mtla, mtlb, mtle, mtlf, mtlg, mtlh, mtli, mtlj, mtlk, mtll, mtlx, mt2, mt2a, mt3, mtacr1, mtap, mtbt1, mtcpl, mterf, mtf1, mth1, mthfc, mthfd, mthfr, mtk1, mtm1, mtmr1, mtmx, mtnrla, mtnrlb, mtp, mtpa, mtr, mtrns, mtrr, mts, mts, mts1, mts1, mts2, mttfl, mtx, mtxn, mu, muc1, muc2, muc3, muc4, muc5, muc5ac, muc5b, muc6, muc8, mul, mum1, mupp1, musk, mut, mvk, mvkl, mvwf, mwfe, mx, mx1, mx2, mxil, mxsl, myb, mybl1, mybl2, mybpc1, mybpc2, mybpc3, mybpcf, mybph, myc, mycl1, mycl2, myclk1, mycn, myd88, myf3, myf4, myf5, myf6, myh1, myh10, myh11, myh12, myh2, myh3, myh4, myh6, myh7, myh8, myh9, myk1, myl, myl1, myl2, myl3, myl4, myl5, mylk, mymy, myo10, myo15, myola, myolc, myold, myole, myo5a, myo6, myo7a, myo9b, myoc, myod1, myog, myp1, myp2, myp3, myr5, mzfl, n33, nab1, nab2, nabc1, nacla, naca, nacao, nacp, nadmr, naga, nage@, naglu, nagr1, naip, namsd, nanta3, nap114, nap2, nap2l, napb, naptb, nars, nat1, nat1, nat2, nb, nb4s, nbat, nbc3, nbccs, nbccs, nbial, nbs, nbs, nbs1, nca, ncad, ncam1, ncan, ncbp, ncc1, ncc2, ncc3, ncc4, ncct, ncf1, ncf2, ncf4, nck, ncl, ncst2, ncx1, ncx2, nd, ndhii, ndn, ndp, ndst1, ndufa1, ndufa2, ndufa5, ndufa6, ndufa7, ndufb8, ndufb9, ndufs1, ndufs2, ndufs4, ndufs7, ndufs8, ndufv1, ndufv2, ndufv3, neb, nec1, nec2, nedd1, nedd2, nedd4, nefh, nefl, negf1, negf2, nell1, nell2, nem1, neol, nep, net, net1, neu, neu, neud4, neurod, neurod2, neurod3, nfl, nfla, nf2, nfatc1, nfatc2, nfatp, nfe1, nfe2, nfe2l1, nfe2l2, nfe2u, nfia, nfib, nfic, nfix, nkfb1, nkfb2, nkfb3, nkfbia, nkfbill, nfrkb, nfya, nfyb, ngal, ngbe, ngfb, ngfg, ngfic, ngfr, ngl, ngn, nhbp, nhcp1, nhcp2, nhe1, nhe3, nhe4, nhe5, nhlh1, nhlh2, nhp2l1, nhs, nid, niddm1, ninj1, nippl, nipsnap1, nipsnap2, nis, nk1r, nkcc1, nkcc2, nkg2, nkg2a, nkg2c, nkg2e, nkg2f, nkhc, nkna, nkna, nknb, nkrp1a, nks1, nksf2, nktr, nkx2a, nkx3.2, nkx3a, nkx6a, nli, nm, nm1, nm23, nmb, nmbr, nmдар1, nmдар2a, nmдар2b, nmдар2c, nmдар2d, nmдара1, nme1, nme2, nme4, nmor1, nmor2, nmsl, nmyc, nnat, nnmt, nno1, nog, nol1, nos1, nos2a, nos2b, nos2c, nos3, not, notch1, notch2, notch3, notch4, nov, nov, nov2, noval, nova3, novp, np, np10, npat, npc, npc1, npd, nph1, nph2, nphl2, nphn, nphp1, nphp2, nphs1, npml, nppa, nppb, npps, npps, npr1, npr2, npr3, nps1, npt1, npt2, nptx2, npy, npy1r, npy2r, npy3r, npy5r, npy6r, nqo2, nramp, nramp1, nramp2, nrap, nras, nrb54, nrcam, nrd1, nrfl, nrfl, nrf2, nrgn, nrip1, nrk2, nrl, nrtn, nru, ns1, nsf, nsp, nspl1, nsrd9, nt4, nt5, nt5, ntcp1, ntcp2, ntf3, ntf4, ntf5, nthl1, ntn, ntn, ntn2l, ntrk1, ntrk2, ntrk3, ntrk4, ntrkr1, ntrkr3, nts, ntt, ntt, nucl, nucb1,

numal, nup214, nup98, nurrl, nvl, nysl, nys2, nysa, oa1, oa2, oa3, oar, oasd, oasl, oat, oat11, oat12, oat13, oatp, oaz, ob, ob10, obf1, obp, obr, oca2, ocm, ocp2, ocrl, ocrl1, oct, oct1, oct1, oct2, oct2, oct3, oct7, octn2, octs3, odc1, odde, odfl, odgl, odod, ofc1, ofc2, ofc3, ofdl, ofe, og12, ogdh, oggl, ogr1, ogs1, ogs2, ohds, ohs, oias, oip1, ok, olf1, olfmf, olfr1, olfr2, omg, omgp, omp, on, op2, opa1, opa2, opa3, opca3, opcm1, opd1, opgl, ophn1, opl1, opn, oppg, oprd1, oprk1, oprm1, oprt, opta2, optb1, oqtl, orld2, orlf1, orc11, orc21, orc41, orc51, orfx, orm1, orm2, orw, osbp, osm, osp, ost, ost48, osx, otc, otfl, otfl2, otfl3, otm, otol, ots, otx1, otx2, ovc, ovcs, ovoll, ox40, oxa11, oxct, oxt, oxtr, ozf, p, p, p1, p15, p16, p167, p28, p2rx3, p2rx4, p2ry1, p2ry2, p2ry4, p2ry7, p2u, p2x3, p2x4, p2y1, p2y2, p2y4, p3, p40phox, p450c11, p450c17, p450c2a, p450c2d, p450c2e, p450sc, p4ha, p4ha1, p4hb, p5cdh, p79r, pa2g4, pab1, pab2, pabp2, pabpl1, pac1, pac1, pacapr, pace, pace4, paep, paf1, paf2, pafah, pafah1b1, pafah1b2, pafah1b3, paga, pah, pahx, pail, pai2, paics, pak1, pak3, palb, pals, pam, pang, pap, papa, papa2, papp, par1, par1, par2, par3, par4, par4, par5, park1, park2, park3, pawr, pax1, pax2, pax3, pax4, pax5, pax6, pax7, pax8, pax9, pbca, pbcr, pbfe, pbgd, pbt, pbx1, pbx2, pbx3, pc, pc1, pc2, pc3, pc3, pca1, pcad, pcap, pcar1, pcbc, pcbd, pcbp1, pcbp2, pcca, pccb, pcdh7, pcdx, pchc, pchc1, pci, pck1, pcl, pclp, pcm1, pcm1, pcmt1, pcna, pcnt, pcolce, pcp, pcp4, pcs, pcsk1, pcsk2, pcsk3, pcsk4, pcsk5, pcsk6, pctk1, pctk3, pcyt1, pdb, pdb2, pdc, pdc, pdc1, pdc2, pddr, pde1a, pde1b, pde1b1, pde3b, pde4a, pde4b, pde4c, pde4d, pde5a, pde6a, pde6b, pde6c, pde6d, pde6g, pde6h, pde7a, pdea, pdea2, pdeb, pdeg, pdes1b, pdgb, pdgfa, pdgfb, pdgfr, pdgfra, pdgfrb, pdha1, pdha2, pdhb, pdj, pdk4, pdnp1, pdnp2, pdnp3, pdr, pds, pds1, pdx1, pdyn, pe1, pe15, pebp2a1, pebp2a3, pecam1, ped, ped, pedf, pee, peg1, peg3, pemp, penk, pent, peo, peo1, peo2, pepa, pepb, pepc, pepd, pepe, pepn, peps, per, per2, peta3, pets1, pex1, pex5, pex6, pex7, pf4, pf4v1, pfas, pfbi, pfc, pfd, pfhb1, pfic1, pfic2, pfkfb1, pfkfb2, pfl, pflm, pfln1, pfln2, pfrx, pga3, pga4, pga5, pgam1, pgam2, pgamm, pgc, pgd, pgf, pgft, pgk1, pgk2, pgka, pgl, pgl1, pgl2, pgm1, pgm2, pgm3, pgm5, pgn, pgp, pgpl, pgr, pgs, pgt, pgy1, pgy3, pha1, pha2, pha2a, pha2b, phap1, phb, phc, phe1a, phe3, phex, phf1, phhi, phhi, phk, phka1, phka2, phkb, phkd, phkg1, phkg2, phl, phl1, phog, phox1, phox2a, php, php1b, php, phyh, pi, pi10, pi3, pi4, pi5, pi6, pi7, pi8, pi9, piga, pigc, pigf, pigh, pigr, pik3c2b, pik3ca, pik3r1, pik4cb, pil, pim1, pin, pin1, pin11, pip, pip5k1b, pir1, pir51, pit, pit1, pitpn, pitx1, pitx2, pitx3, pjs, pk1, pk120, pk3,



ptprl2, ptpm, ptpn, ptpo, ptp, ptpz1, ptp, pts, pti, pti, pti, pti, pujo, pum, pur1, pur1, pura, pvalb, pvr, pvr1, pvr2, pvr1, pvr2, pvs, pvt1, pwcr, pwp2, pwp2h, pws, pxaal, pxe, pxel, pxf, pxmp1, pxmp11, pxmp3, pxr1, pyer1, pyes, pygb, pygl, pygm, pyk2, pyst1, pyst2, pzp, qars, qdpr, qin, qm, qpc, qpr, rab, rab1, rab13, rab1a, rab21, rab3a, rab3b, rab4, rab5, rab5a, rab6, rab7, rabgd1a, rabgdib, rabggta, rabggtb, rabif, rac2, rac3, rad1, rad17, rad23a, rad23b, rad51a, rad51c, rad51d, rad51l3, rad52, rad54, rad6a, rad6b, raf1, rafal, rag1, rag2, rage, rala, ralb, ralgs, ramp, ranbp211, ranbp3, rao, rap1a, rap1b, rap1ga1, rap1gds1, rap2a, rap74, rapsn, rara, rarb, rarg, rars, rasal, rasa2, rasgfr3, rask2, rb1, rbbp2, rbbp5, rbbp6, rbl1, rbl2, rbm1, rbm2, rbm3, rbmy1a1, rbp1, rbp2, rbp3, rbp4, rbp5, rbp56, rbp6, rbq3, rbtn1, rbtn11, rbtn12, rcal, rcac, rcc1, rccp1, rccp2, rcd1, rcd2, rcdp1, rcn1, rcn2, rcp, rcv1, rd, rdbp, rdc7, rdp, rdpa, rdrc, rds, rdt, rdx, reca, reccl, recql, red1, red2, reg, reg1a, regl, rel, rela, reln, ren, renbp, rens1, rent1, rep8, req, ret, rev3, rev31, rfc1, rfc2, rfc3, rfc4, rfc5, rfp, rfx1, rfx2, rfx5, rfxank, rfxap, rgcl, rgr, rgs, rgs1, rgs14, rgs16, rgs2, rgs2, rgs3, rgs5, rh50a, rhag, rhbd1, rhc, rhce, rhd, rheb2, rho, rho7, rhogap2, rhogap3, rhoh12, rhoh6, rhoh9, rhok, rhom1, rhom2, rhom3, rieg1, rieg2, rige, rigui, ring1, ring10, ring11, ring12, ring3, ring31, ring4, ring5, ring6, ring7, rip, rip140, riz, rk, rl, rlb1, rlf, rln1, rln2, rmch1, rmd1, rmrp, rmrpr, m5s1, mase1, mase2, mase3, mase4, mase5, mase6, mase1, mase1, me1, mfl, mfl3, mfl4, mfl5, mnh, mnep, mpulz, mnr1, mnr2, mnr3, mnr4, mnr5, mns1, mns2, mns3, mns4, mns4, mns4i, mntmi, mu1, mu15a, mu17a, mu17b, mula, mu2, mu3, ro52, rom1, romk1, ron, ror1, rora, rorb, rorc, rorg, ros1, rosp1, rox, rp1, rp10, rp105, rp11, rp12, rp13, rp14, rp15, rp17, rp18, rp19, rp2, rp22, rp24, rp25, rp3, rp4, rp6, rp7, rp9, rpa1, rpa2, rpa3, rpd311, rpe, rpe65, rpl19, rpl22, rpl23a, rpl231, rpl29, rpl30, rpl35a, rpl36a, rpl7a, rpsm12, rpn1, rpn2, rpol2, rps11, rps14, rps17, rps17a, rps17b, rps17l1, rps17l2, rps18, rps20a, rps20b, rps24, rps25, rps3, rps4x, rps4y, rps6, rps6ka1, rps6ka2, rps6ka3, rps8, rpsm12, rptpm, rpul, rpx, rrad, rras, rrbp1, rreb1, rrm1, rrm2, rrp, rrp22, rs1, rs1, rsclal, rsk1, rsk2, rsk3, rsn, rss, rsts, rsul, rt6, rtefl, rtkn, rtn1, rtn2, rts, rts, rtt, rws, rxra, rxrb, rxrg, ryr1, ryr2, ryr3, rzrb, rzrg, sl00a1, sl00a10, sl00a11, sl00a12, sl00a13, sl00a2, sl00a3, sl00a4, sl00a5, sl00a6, sl00a7, sl00a8, sl00a9, sl00b, sl00d, sl00e, sl00l, sl00p, sl52, s4, s7, saal, saa2, saa4, sacs, safb, sag, sah, sahh, sail, sakap84, sall1, sall2, sams1, sams2, sap, sap1, sap1, sap2, sap62, sar, sar1, sar2, sard, sas, sat, satb1, satt, sbma, sc, scl1, scl5dl, scal, scal0, sca2, sca2, sca3, sca4,

sca5, sca6, sca7, sca8, sca8, scar, scca1, scca2, sccd, scd, sceh, scg1, scg2, scg3, schad, scida, scidx, scidx1, scl, sclc1, scll, scn, scn1a, scn1b, scn2a, scn2a1, scn2a2, scn2b, scn3a, scn4a, scn5a, scn6a, scn8a, scnna1, scnna1b, scnna1d, scnna1g, scot, scp, scp1, scp2, scpn, scra1, scra1, scs, sctr, scya1, scya11, scya13, scya14, scya15, scya16, scya19, scya2, scya21, scya22, scya24, scya25, scya3, scya311, scya4, scya5, scya7, scya8, scyb5, scyb6, scyd1, sczd1, sczd2, sczd3, sczd4, sczd5, sczd6, sczd7, sczd8, sdc1, sdc2, sdc4, sdf1, sdf2, sdh1, sdh2, sdha, sdhb, sdhc, sdhd, sdhf, sds22, sdty3, sdys, se, sea, sec1311, sec13r, sec141, sec7, sed1, sedt, sef2, sel11, sele, sell, selp, selplg, sema3f, sema4, sema5, semg, semg1, semg2, sen1, sep, sepp1, serca1, serca3, serk1, ses1, set, sex, sf, sfl, sfa1, sfd, sfmd, sfrs1, sfrs2, sfrs7, sftb3, sftp1, sftp2, sftp4, sftpa1, sftpa2, sftpb, sftpc, sftpd, sgb, sgca, sgcb, sgcd, sgcg, sgd, sgk, sglt1, sglt2, sgm1, sgne1, sgp2, sgpa, sgsh, sh2d1a, sh3bp2, sh3d1a, sh3gbr, sh3p17, shb, shbg, shc1, shcl1, shfd1, shfd2, shfm1, shfm2, shfm3, shh, ship, shmt1, shmt2, shoc2, shot, shox, shox2, shps1, shs, shsf1, si, siah1, siah2, siasd, siat1, siat4, siat4c, siat8, sids, sil, silv, sim1, sim2, sipa1, sis, siv, six1, six5, sja, sjs, ski, ski2, ski2w, skiv21, skp1a, skp1b, skp2, sla, slap, slbp, slc, slc10a1, slc10a2, slc12a1, slc12a2, slc12a3, slc14a1, slc14a2, slc15a1, slc16a1, slc16a2, slc17a1, slc17a2, slc18a1, slc18a2, slc18a3, slc19a1, slc1a1, slc1a2, slc1a3, slc1a4, slc1a5, slc20a1, slc20a2, slc20a3, slc21a2, slc21a3, slc22a1, slc22a2, slc22a5, slc2a1, slc2a2, slc2a3, slc2a4, slc2a5, slc2c, slc3a1, slc4a1, slc4a2, slc4a6, slc5a1, slc5a2, slc5a3, slc5a5, slc6a1, slc6a10, slc6a12, slc6a2, slc6a3, slc6a4, slc6a6, slc6a8, slc6a9, slc7a1, slc7a2, slc7a4, slc7a5, slc7a7, slc8a1, slc8a2, slc9a1, slc9a2, slc9a3, slc9a4, slc9a5, sld, sle1, sleb1, slim1, sln, slo, slos, slp76, sls, slug, sm1, sm22, sma4, smad1, smad1, smad2, smad3, smad4, smad5, smad6, smad7, smad9, smal, smam1, smarca1, smarca2, smarca3, smarca5, smarcb1, smax2, smc1, smcc, smcr, smcx, smcy, sml1, smn, smn1, smn2, smnr, smo, smoh, smpd1, sms, smt3, smt3h1, smtn, smubp2, sn, snap25, snat, snca, sncb, sncg, snf2h, snf211, snf212, snf213, snf5, snl, snn, snrp70, snrpa, snrpe, snrpn, snt1, snt2b1, snt2b2, sntb1, sntl, snx, soat, sod1, sod2, sod3, solh, son, sord, sor11, sos1, sos2, sox1, sox10, sox11, sox2, sox20, sox22, sox3, sox4, sox9, sp1, sp1, sp3, sp3, sp4, spa1, spag1, spag4, spam1, sparc, spat, spbp, spch1, spd, spf30, spg3a, spg4, spg5a, spg6, spg7, spg8, spg9, spgp, spgyla, sph2, spi1, spink1, spk, spmd, spn, spp1, spp2, sppm, spr, sprk, sprr1a, sprr1b, sprr2a, sprr2b, sprr2c, sprr3, sps1, spsma, spta1, sptan1, sptb, sptbn1, sra1, sra2, src, src1, src1, src2, srd5a1,



tric5, trio, trip1, trip14, trip6, trk, trk1, trka, trkb, trkc, trke, trl1, trl2, trm1, trm1, trm2, trma,  
 trmi1, trmi2, trn, trnl, tro, trp1, trp1, trp2, trp3, trpc1, trpm2, trpo, trps1, trps2, trq1, trr, trr3,  
 trrap, trsp, trt1, trt2, trv1, trv2, trv3, trv4, trv5, try1, try2, ts, ts13, ts546, tsbn51, tsc, tsc1,  
 tsc2, tsd, tsel, tsg101, tsg7, tshb, tshr, tsix, tsp3, tspy, tssc3, tst1, tst1, tsta3, tsy, ttc1, ttc3, ttf,  
 ttf1, ttf2, ttg2, ttim1, ttn, ttp, ttp1, ttpa, ttr, tuba3, tuba1, tubb, tufm, tuft1, tulp1, tuple1, tw,  
 tweak, twik1, twist, txgp11, txk, txn, txnr, txnrd1, tyh, tyk1, tyk2, tyk3, tyms, tyr, tyrl, tyro3,  
 tyrp1, tyrp2, tys, u17hg, ulrnp, u22hg, u2af1, u2af1rs1, u2af1rs2, u2af1rs3, uba52, ubb, ubc,  
 ubc4, ubc7, ubc8, ubch2, ubcl, ube1, ube2, ube2a, ube2b, ube2e2, ube2g, ube2g2, ube2h,  
 ube2i, ube2i1, ube2v1, ube3a, ubh1, ubid4, ubl1, uch11, ucn, ucp1, ucp2, ucp3, udpgdh, uev1,  
 ufd11, ufs, ugalt, ugb, ugcg, ugdh, ugn, ugp1, ugp2, ugpp2, ugt1, ugt1a1, ugt2b11, ugt2b15,  
 ugt2b17, ugt2b4, ugt2b7, ugt2b8, ugt2b9, ugt1, uhg, uhx1, ukhc, umod, umph2, umpk, umps,  
 unc18, unc18b, und, ung, unr, uox, up, upk1b, ups, uqbp, uqcrb, uqcrc1, uqcrc2, uqcrcs1,  
 uqor1, uqor13, uqor22, urk, urkr, uroc, urod, uros, usf1, usf2, ush1, ush1a, ush1b, ush1c,  
 ush1d, ush1e, ush1f, ush2a, ush3, usp11, usp5, usp7, usp9x, usp9y, ut1, ut2, ute, utr, utrn,  
 utx, uty, uv20, uv24, uvo, vacht, vacm1, vamp1, vamp2, vars1, vasp, vat1, vat2, vav, vav1,  
 vav2, vbch, vbpl, vcam1, vcf, vcl, vcp, vdac1, vdac2, vdd1, vdi, vdr, vegf, vegfb, vegfd,  
 vegfr3, vgf, vgl, vgr1, vhl, vhr, vill, vil2, vim, vip, vipr1, vipr2, vis1, vla1, vla5a, vlacs,  
 vlcad, vldlr, vmat1, vmcm, vmd1, vmd2, vnra, vnt, vp, vpp1, vpp3, vpreb1, vpreb2, vrf, vrk1,  
 vrk2, vrnf, vrni, vsn11, vtn, vwf, vws, waf1, wars, was, wbs, wd1, wdr2, weel, wfes, wfs,  
 wfs1, wgn1, whcr, wi, wisp1, wisp2, wisp3, wnd, wnt1, wnt10b, wnt13, wnt14, wnt15, wnt2,  
 wnt3, wnt5a, wnt7a, wnt7b, wnt8b, wrb, wrn, ws1, ws2a, ws2b, ws4, wsn, wss, wss, wt1,  
 wt2, wt3, wt4, wt5, wts, wtsl, wws, x11, xbp1, xbp2, xce, xdh, xe169, xe7, xe7y, xg, xgr,  
 xh2, xiap, xic, xist, xk, xla, xla2, xlp, xlpd, xlr1, xm, xpa, xpb, xpc, xpcc, xpct, xpf, xpf,  
 xpg, xpmc2h, xpnpep2, xpo1, xrcc1, xrcc2, xrcc3, xrcc4, xrcc5, xrcc9, xrs, xs, xwnt2, yb1,  
 yes1, ykl40, yll, yrrm1, yt, ywha1, ywhab, ywhah, ywhaz, yy1, zac, zag, zan, zap70, zf87,  
 zfm1, zfp3, zfp36, zfp37, zfx, zfy, zic1, zic2, zic3, zipk, znf1, znf10, znf117, znf11a, znf11b,  
 znf12, znf121, znf123, znf124, znf125, znf126, znf13, znf14, znf141, znf144, znf146, znf147,  
 znf157, znf16, znf160, znf162, znf163, znf165, znf169, znf173, znf179, znf18, znf189, znf19,  
 znf192, znf193, znf195, znf198, znf2, znf20, znf200, znf204, znf217, znf22, znf23, znf24,  
 znf25, znf26, znf27, znf29, znf3, znf32, znf34, znf35, znf36, znf38, znf4, znf40, znf41,

znf42, znf44, znf45, znf46, znf5, znf6, znf69, znf7, znf70, znf71, znf72, znf73, znf74, znf75, znf75a, znf75c, znf76, znf77, znf79, znf8, znf80, znf81, znf83, znf9, znfc150, znfc25, znfx, znt3, znt4, zp3a, zp3b, zpk, zws1, and zyx.

[0065] Furthermore, genes from bacteria, plants, yeast, and mammals (e.g., mice) can be used with the present invention. Examples of *E. coli* genes include: aarF, aas, aat, abpS, abs, accA, accB, accC, accD, acd, aceA, aceB, aceE, aceF, aceK, ackA, ackB, acnA, acnB, acpD, acpP, acpS, acpX, acrA, acrB, acrC, acrD, acrE, acrF, acrR, acs, ada, add, adhB, adhC, adhE, adhR, adiA, adiY, adk, aegA, aer, aes, agaA, agaB, agaC, agaD, agaI, agaR, agaS, agaV, agaW, agaZ, agp, ahpC, ahpF, aidB, ais, alaS, alaT, alaU, alaV, alaW, alaX, aldA, aldB, aldH, alkA, alkB, alpA, alr, alsA, alsB, alsC, alsE, alsK, alx, amiA, amiB, amn, ampC, ampD, ampE, ampG, ampH, amtB, amyA, ansA, ansB, apaG, apaH, aphA, appA, appB, appC, appY, apt, aqpZ, araA, araB, araC, araD, araE, araF, araG, araH, araJ, arcA, arcB, argA, argB, argC, argD, argE, argF, argG, argH, argI, argM, argP, argQ, argR, argS, argT, argU, argV, argW, argX, argY, argZ, aroA, aroB, aroC, aroD, aroE, aroF, aroG, aroH, aroI, aroK, aroL, aroM, aroP, aroT, arsB, arsC, arsR, artI, artJ, artM, artP, artQ, ascB, ascF, ascG, asd, aslA, aslB, asmA, asnA, asnB, asnC, asnS, asnT, asnU, asnV, asnW, aspA, aspC, aspS, aspT, aspU, aspV, asr, asu, atoA, atoB, atoC, atoD, atoS, atpA, atpB, atpC, atpD, atpE, atpF, atpG, atpH, atpI, avtA, azaA, azaB, azl, bacA, baeR, baeS, barA, basR, basS, bax, bcp, bcr, betA, betB, betI, betT, bfd, bfm, bfr, bglA, bglB, bglF, bglG, bglJ, bglT, bglX, bioA, bioB, bioC, bioD, bioF, bioH, bioP, bipA, birA, bisC, bisZ, blc, bolA, brnQ, brnR, brnS-??, brnT, btuB, btuC, btuD, btuE, btuR, bymA, cadA, cadB, cadC, cafA, caiA, caiB, caiC, caiD, caiE, caiF, caiT, calA, calC, calD, can, carA, carB, cbl, cbpA, cbt, cca, ccmA, ccmB, ccmC, ccmD, ccmE, ccmF, ccmG, ccmH, cdd, cde, cdh, cdsA, cdsS, cedA, celA, celB, celC, celD, celF, cfa, cfcA, chaA, chaB, chaC, cheA, cheB, cheR, cheW, cheY, cheZ, chpA, chpB, chpR, chpS, cirA, citA, citB, cld, clpA, clpB, clpP, clpX, cls, cmk, cmlA, cmr, cmtA, cmtB, coaA, cobS, cobT, cobU, codA, codB, cof, cog?, corA, cpdA, cpdB, cpsA, cpsB, cpsC, cpsD, cpsE, cpsF, cpsG, cpxA, cpxB, cpxP, cpxR, crcA, crcB, creA, creB, creC, creD, crg, srl, crp, crr, csdA, csgA, csgB, csgD, csgE, csgF, csgG, csiA, csiB, csiC, csiD, csiE, csiF, cspA, cspB, cspC, cspD, cspE, cspG, csrA, csrB, cstA, cstC, cup, cutA, cutC, cutE, cutF, cvaA(ColV), cvaB(ColV), cvaC(ColV), cvi(ColV), cvpA, cxm, cyaA, cybB, cybC, cycA, cydA, cydB,



cydC, cydD, cynR, cynS, cynT, cynX, cyoA, cyoB, cyoC, cyoD, cyoE, cysA, cysB, cysC, cysD, cysE, cysG, cysH, cysI, cysJ, cysK, cysM, cysN, cysP, cysQ, cysS, cysT, cysU, cysW, cysX?, cysZ?, cytR, dacA, dacB, dacC, dacD, dadA, dadB, dadQ, dadX, dam, dapA, dapB, dapD, dapE, dapF, dbpA, dcd, dcm, dcp, dcrB, dctA, dctB, dcuA, dcuB, dcuC, ddlA, ddlB, ddpA, ddpB, ddpC, ddpD, ddpF, ddpX, deaD, dedA, dedD, def, degP, degQ, degS, del, deoA, deoB, deoC, deoD, deoR, dfp, dgd, dgkA, dgkR, dgoA, dgoD, dgoK, dgoR, dgoT, dgsA, dgt, dicA, dicB, dicC, dicF, dinB, dinD, dinF, dinG, dinI, dinY, dipZ, djlA, dksA, dld, dmsA, dmsB, dmsC, dnaA, dnaB, dnaC, dnaE, dnaG, dnaI, dnaJ, dnaK, dnaL, dnaN, dnaQ, dnaT, dnaX, dppA, dppB, dppC, dppD, dppF, dppG, dps, dsbA, dsbB, dsbC, dsbG, dsdA, dsdC, dsdX, dsrA, dsrB, dut, dvl, dxs, ebgA, ebgB, ebgC, ebgR, ecfA, eco, ecpD, eda, edd, efp, emrA, emrB, emrD, emrE, endA, eno, entA, entB, entC, entD, entE, entF, envN-??, envP, envQ, envR, envT, envY, envZ, epd, EppA, minigene, EppB, minigene, EppC, minigene, EppD, minigene, EppE, minigene, EppG, minigene, EppH, minigene, era, esp, evgA, evgS, exbB, exbC, exbD, expA, exuR, exuT, fabA, fabB, fabD, fabF, fabG, fabH, fabI, fabZ, fadA, fadB, fadD, fadE, fadH, fadL, fadR, farR, fatA, fbaA, fbaB, fbp, fcl, fcsA, fdhD, fdhE, fdhF, fdnG, fdnH, fdnI, fdoG, fdoH, fdoI, fdrA, fdx, feaB, feaR, fecA, fecB, fecC, fecD, fecE, fecI, fecR, feoA, feoB, fepA, fepB, fepC, fepD, fepE, fepG, fes, fexB, ffh, ffs, fhlA, fhlB, fhuA, fhuB, fhuC, fhuD, fhuE, fhuF, fic, fimA, fimB, fimC, fimD, fimE, fimF, fimG, fimH, fimI, fipB, fipC, fis, fiu, fixA, fixB, fixC, fixX, fklB, fkpA, fldA, flgA, flgB, flgC, flgD, flgE, flgF, flgG, flgH, flgI, flgJ, flgK, flgL, flgM, flgN, flhA, flhB, flhC, flhD, fliA, fliC, fliD, fliE, fliF, fliG, fliH, fliI, fliJ, fliK, fliL, fliM, fliN, fliO, fliP, fliQ, fliR, fliS, fliT, fliY, fliZ, flk, flu, fnt, fnr, focA, focB, folA, folC, folD, folE, folK, folP, folX, fpr, frdA, frdB, frdC, frdD, frr, fruA, fruB, fruK, fruR, fsr, ftn, ftsA, ftsE, ftsI, ftsJ, ftsK, ftsL, ftsN, ftsQ, ftsW, ftsX, ftsY, ftsZ, fucA, fucI, fucK, fucO, fucP, fucR, fumA, fumB, fumC, fur, fusA, fusB, gabC-??, gabD, gabP, gabT, gadA, gadB, gadR, galE, galF, galK, galM, galP, galR, galS, galT, galU, gapA, gapC, garA, garB, gatA, gatB, gatC, gatD, gatR, gatY, gatZ, gcd, gcl, gcpE, gcvA, gcvH, gcvP, gcvR, gcvT, gdhA, gef, ggt, gidA, gidB, gip, glcB, glcC, glcD, glcE, glcG, gldA, glf, glgA, glgB, glgC, glgP, glgS, glgX, glk, glmM, glmS, glmU, glmX, glnA, glnB, glnD, glnE, glnG, glnH, glnK, glnL, glnP, glnQ, glnR, glnS, glnT, glnU, glnV, glnW, glnX, gloA, glpA, glpB, glpC, glpD, glpE, glpF, glpG, glpK, glpQ, glpR,

glpT, glpX, gltA, gltB, gltD, gltE, gltF, gltH, gltJ, gltK, gltL, gltM, gltP, gltR, gltS, gltT, gltU, gltV, gltW, gltX, glyA, glyQ, glyS, glyT, glyU, glyV, glyW, glyX, glyY, gmd, gmk, gmm, gnd, gntK, gntP, gntR, gntS, gntT, gntU, gntV, goaG, gor, gph, gpmA, gpp, gprA, gprB, gpsA, gpt, greA, greB, groL, groS, grpE, grxA, grxB, grxC, gshA, gshB, gsk, gsp, gsp\*, gst, guaA, guaB, guaC, gurB, gurC, gutM, gutQ, gyrA, gyrB, hcaB, hcaC, hcaD, hcaE, hcaF, hcaR, hcaT, hdeA, hdeB, hdeD, hdhA, helD, hemaA, hemB, hemC, hemD, hemE, hemF, hemG, hemH, hemK, hemL, hemM, hemX, hemY, hepA, het, hflB, hflC, hflK, hflX, hfq, hha, hipA, hipB, hisA, hisB, hisC, hisD, hisF, hisG, hisH, hisI, hisJ, hisM, hisP, hisQ, hisR, hisS, hlpA, hlyE, hmp, hns, holA, holB, holC, holD, holE, hopB, hopC, hopD, hpt, hrpA, hrpB, hrsA, hscA, hscB, hsdM, hsdR, hsdS, hslC, hslD?, hslE-H, hslJ, hslK, hslL-N, hslO-R, hslU, hslV, hslW, htgA, htpG, htpX, htrB, htrC, htrE, htrL, hupA, hupB, hyaA, hyaB, hyaC, hyaD, hyaE, hyaF, hybA, hybB, hybC, hybD, hybE, hybF, hybG, hycA, hycB, hycC, hycD, hycE, hycF, hycG, hycH, hycI, hydA, hydG, hydH, hydN, hyfA, hyfB, hyfC, hyfD, hyfE, hyfF, hyfG, hyfH, hyfI, hyfJ, hyfR, hypA, hypB, hypC, hypD, hypE, hypF, iadA, iap, ibpA, ibpB, icd, iclR, ihfA, ihfB, ileR, ileS, ileT, ileU, ileV, ileX, ileY, ilvA, ilvB, ilvC, ilvD, ilvE, ilvF, ilvG, ilvH, ilvI, ilvJ-??, ilvM, ilvN, ilvR, ilvU, ilvY, imp, inaA, inaR?, infA, infB, infC, inm, insA(IS1), intA, isb(IS1), isfA, ispA, ispB, KanR, katE, katG, kba, kbl, kch, kdgK, kdgR, kdgT, kdpA, kdpB, kdpC, kdpD, kdpE, kdpF, kdsA, kdsB, kdtA, kdtB, kefB, kefC, kgtP, ksgA, ksgB, ksgC, ksgD, lacA, lacI, lacY, lacZ, lamB, lar, ldcC, ldhA, lepA, lepB, leuA, leuB, leuC, leuD, leuJ, leuO, leuP, leuQ, leuR, leuS, leuT, leuU, leuV, leuW, leuX, leuY, leuZ, lev, lexA, lgt, lhr, ligA, ligT, linB, lipA, lipB, lit, livF, livG, livH, livJ, livK, livM, lldD, lldP, lldR, lolA, lon, lpcA, lpcB, lpd, lplA, lpp, lpxA, lpxB, lpxC, lpxD, lpxK, lrb, lrhA, lrp, lrs-??, lspA, lysA, lysC, lysP, lysQ, lysR, lysS, lysT, lysU, lysV, lysW, lysX, lysY, lysZ, lytA, lytB, lyx, maa, mac, mae, mafA, mafB, malE, malF, malG, malI, malK, malM, malP, malQ, malS, malT, malX, malY, malZ, manA, manC, manX, manY, manZ, map, marA, marB, marR, mbrB, mcrA, mcrB, mcrC, mcrD, mdaB, mdh, mdoB, mdoG, mdoH, meb, melA, melB, melR, menA, menB, menC, menD, menE, menF, mepA, mesJ, metA, metB, metC, metD, metE, metF, metG, metH, metJ, metK, metL, metR, metT, metU, metV, metW, metY, metZ, mfd, mglA, mglB, mglC, mglR, mgsA, mgtA, mhpA, mhpB, mhpC, mhpD, mhpE, mhpF, mhpR, miaA, miaD, micF, minC, minD, minE, mioC,

mltA, mltB, mltC, mltD, mmrA(rhlB?), mng, mntA, moaA, moaB, moaC, moaD, moaE, mobA, mobB, moc, modA, modB, modC, modE, modF, moeA, moeB, mog, molR, motA, motB, mpl, mppA, mprA, mraA--?, mraY, mrcA, mrcB, mrdA, mrdB, mreB, mreC, mreD, mrp, mrr, msbA, msbB, mscL, msrA, msyB, mtg, mtgA, mtlA, mtlD, mtlR, mtr, mttA, mttB, mttC, mukB, mukE, mukF, mul, murA, murB, murC, murD, murE, murF, murG, murH, murI, mutG(putative), mutH, mutL, mutM, mutS, mutT, mutY, nac, nadA, nadB, nadC, nadE, nagA, nagB, nagC, nagD, nagE, nalB, nalD, nanA, nanE, nanK, nanR, nanT, napA, napB, napC, napD, napF, napG, napH, narG, narH, narI, narJ, narK, narL, narP, narQ, narU, narV, narW, narX, narY, narZ, ndh, ndk, neaB, nei, nemA, nfi, nfnA, nfnB, nfo, nfrA, nfrB, nfrD, nfsA, nhaA, nhaB, nhaR, nikA, nikB, nikC, nikD, nkeE, nirB, nirC, nirD, nlpA, nlpB, nlpC, nlpD, nmpC(qsr'), non, npr, nrdA, nrdB, nrdD, nrdE, nrdF, nrdG, nrfA, nrfB, nrfC, nrfD, nrfE, nrfF, nrfG, nth, ntpA, nuoA, nuoB, nuoC, nuoE, nuoF, nuoG, nuoH, nuoI, nuoJ, nuoK, nuoL, nuoM, nuoN, nupC, nupG, nusA, nusB, nusG, nuvA, nuvC, ogrK, ogt, ompA, ompC, ompF, ompG, ompR, ompT, ompX, oppA, oppB, oppC, oppD, oppE, oppF, opr, ops, oraA, ordL, orf-23(purB, reg)orf195(nikA-reg), orn, osmB, osmC, osmE, osmY, otsA, otsB, oxyR, oxyS, pabA, pabB, pabC, pac, pal, panB, panC, panD, panF, parC, parE, pat, pbpG, pck, pcm, pcnB, pdhR, pdxA, pdxB, pdxH, pdxJ, pdxK, pdxL, pdxY, pepA, pepD, pepE, pepN, pepP, pepQ, pepT, pfkA, pfkB, pflA, pflB, pflC, pflD, pfs, pgi, pgk, pgl, pgm, pgpA, pgpB, pgsA, pheA, pheP, pheS, pheT, pheU, pheV, phnC, phnD, phnE, phnF, phnG, phnH, phnI, phnJ, phnK, phnL, phnM, phnN, phnO, phnP, phoA, phoB, phoE, phoH, phoP, phoQ, phoR, phoU, phrB, phxB, pin, pioO, pit, pldA, pldB, plsB, plsC, plsX, pmbA, pncA, pncB, pnp, pntA, pntB, pnuC, poaR, polA, polB, popD, potA, potB, potC, potD, potE, potF, potG, potH, potI, poxA, poxB, ppa, ppc, pphA, pphB, ppiA, ppiB, ppiC, ppk, pppA, pps, ppx, pqiA, pqiB, pqqL, pqqM, prc, prfA, prfB, prfC, priA, priB, priC, prlC, prlZ, prmA, prmB, proA, proB, proC, proK, proL, proM, proP, proQ, proS, proT, proV, proW, proX, prpA, prpC, prpR, prr, prs, psd, psiF, pspA, pspB, pspC, pspE, pspF, pssA, pssR, pstA, pstB, pstC, pstS, psu, pta, pth, ptrA, ptrB, ptsG, ptsH, ptsI, ptsN"-", ptsP, purA, purB, purC, purD, purE, purF, purH, purK, purL, purM, purN, purP, purR, purT, purU, pus, putA, putP, pykA, pykF, pyrB, pyrC, pyrD, pyrE, pyrF, pyrG, pyrH, pyrI, qmeC, qmeD, qmeE, qor, queA, racC, racR, radA, radC, ranA, rarD, ras, rbfA, rbn, rbsA, rbsB, rbsC, rbsD, rbsK, rbsR, rcsA, rcsB, rcsC,

rcsF, rdgA, rdgB, recA, recB, recC, recD, recE, recF, recG, recJ, recN, recO, recQ, recR, recT, relA, relB, relE, relF, relX, rep, rer, rfaB, rfaC, rfaD, rfaF, rfaG, rfaH, rfaI, rfaJ, rfaK, rfaL, rfaP, rfaQ, rfaS, rfaY, rfaZ, rfbA, rfbB, rfbC, rfbD, rfbX, rfc, rfe, rffA, rffC, rffD, rffE, rffG, rffH, rffM, rffT, rhaA, rhaB, rhaD, rhaR, rhaS, rhaT, rhlB, rhlE, rho, ribA, ribB, ribC, ribD, ribE, ribF, ridA, ridB, rimB, rimC, rimD, rimE, rimG, rimH, rimI, rimJ, rimK, rimL, rimM, rit, rlpA, rlpB, rluA, rluC, rluD, rmf, rna, rnb, rnc, rnd, rne, rnhA, rnhB, mnk, mpA, mpB, mr, rnt, rob, rorB, rpe, rph, rpiA, rpiB, rpiR, rplA, rplB, rplC, rplD, rplE, rplF, rplI, rplJ, rplK, rplL, rplM, rplN, rplO, rplP, rplQ, rplR, rplS, rplT, rplU, rplV, rplW, rplX, rplY, rpmA, rpmB, rpmC, rpmD, rpmE, rpmF, rpmG, rpmH, rpmI, rpmJ, rpoA, rpoB, rpoC, rpoD, rpoE, rpoH, rpoN, rpoS, rpoZ, rpsA, rpsB, rpsC, rpsD, rpsE, rpsF, rpsG, rpsH, rpsI, rpsJ, rpsK, rpsL, rpsM, rpsN, rpsO, rpsP, rpsQ, rpsR, rpsS, rpsT, rpsU, rrfA, rrfB, rrfC, rrfD, rrfE, rrfF, rrfG, rrfH, rrlA, rrlB, rrlC, rrlD, rrlE, rrlG, rrlH, rrmA, rrsA, rrsB, rrsC, rrsD, rrsE, rrsG, rrsH, rsd, rseA, rseB, rseC, rspA, rspB, rssA, rssB, rsuA, rtcA, rtcB, rtcR, rtn, rus(qsr'), ruvA, ruvB, ruvC, sad, sanA, sapA, sapB, sapC, sapD, sapF, sbaA, sbcB, sbcC, sbcD, sbmA, sbmC(gyrI), sbp, sdaA, sdaB, sdaC, sdhA, sdhB, sdhC, sdhD, sdiA, sds, secA, secB, secD, secE, secF, secG, secY, selA, selB, selC, selD, semA, seqA, serA, serB, serC, serR-??, serS, serT, serU, serV, serW, serX, sfa, sfcA, sfiC, sfsA, sfsB, shiA, sipC, sipD, sir, sixA, sloB, slp, slr, slt, slyD, slyX, smp, smtA, sodA, sodB, sodC, sohA, sohB, solA, soxR, soxS, speA, speB, speC, speD, speE, speF, speG, spf, spoT, sppA, spr, srlA, srlB, srlD, srlE, srlR, srmB, smA, ssaE, ssaG, ssaH, ssb, sseA, sseB, sspA, sspB, ssrA, ssrS, ssyA, ssyD-??, stfZ, stkA, stkB, stkC, stkD, stpA, strC, strM, stsA, sucA, sucB, sucC, sucD, sufl, sugE, suhA, suhB, sula, supQ, surA, surE, syd, tabC, tag, talA, talB, tanA, tanB, tap, tar, tas, tauA, tauB, tauC, tauD, tbpA, tdcA, tdcB, tdcC, tdcD, tdcE, tdcF, tdcG, tdcR, tdh, tdi-??, tdk, tehA, tehB, tesA, tesB, tgt, thdA, thdC, thdD, thiB?, thiC, thiD, thiE, thiF, thiG, thiH, thiI, thiJ, thiK, thiL, thiM, thrA, thrB, thrC, thrS, thrT, thrU, thrV, thrW, thyA, tig, tktA, tktB, tldD, tlnA, tmk, tnaA, tnaB, tnaC, tnm, tol-orf1, tol-orf2, tolA, tolB, tolC, tolD, tolE, tolI, tolJ, tolM, tolQ, tolR, tonB, topA, topB, torA, torC, torD, torR, torS, torT, tpiA, tpr, tpx, treA, treB, treC, treF, treR, trg, trkA, trkD, trkG, trkH, trmA, trmB, trmC, trmD, trmE, trmF, trmH, trmU, trnA, trpA, trpB, trpC, trpD, trpE, trpR, trpS, trpT, truA, truB, trxA, trxB, trxC, tsA, tsf, tsmA, tsr, tsx, ttdA, ttdB, ttk, tufA, tufB, tus, tynA, tyrA, tyrB, tyrP, tyrR, tyrS, tyrT, tyrU, tyrV, ubiA,

ubiB, ubiC, ubiD, ubiE, ubiF, ubiG, ubiH, ubiX, ucpA[], udk, udp, ugpA, ugpB, ugpC, ugpE, ugpQ, uhpA, uhpB, uhpC, uhpT, uidA, uidB, uidR, umuC, umuD, ung, upp, uppS, ups, uraA, usg-1, ushA, uspA, uup, uvh, uvrA, uvrB, uvrC, uvrD, uvs, uxaA, uxaB, uxaC, uxuA, uxuB, uxuR, valS, valT, valU, valV, valW, valX, valY, valZ, vsr, wrbA, xapA, xapB, xapR, xasA, xerC, xerD, xni, xseA, xseB, xthA, xylA, xylB, xylE, xylF, xylG, xylH, xylR, ycca, yhhP, yihG, yjaB, fl47, yjaD, yohF, yqiE, yrfE, zipA, zntA, znuA, znuB, znuC, zur, and zwf.

[0066] Examples of mouse genes include: Ilr1, Ilr2, Gas10, Tnp1, Inhbb, Inha, Creb1, Mpmv34, Acrd, Acrg, Il10, Otf1, Rab11b-r, Abll, ald, Amh-rs1, Bcl2B, Cchl1a3, Ccnb1-rs2, Gpcr16, Htr5b, Idd5, Igfbp2, Igfbp5, Il8rb, Kras2-rs1, Mov7, Mpmv6, Mpmv16, Mpmv22, Mpmv25, Mpmv29, Mpmv42, Mtv7, Mtv27, Mtv39, Oprk1, Otf3-rs1, Otf8, Otf11-rs1, Ptgs2, Ren1, Ren2, Ril3, Sxv, Taz4-rs1, Tgfb2, Wnt6, Xmmv6, Xmmv9, Xmmv36, Xmmv61, Xmmv74, Xmv21, Xmv32, Xmv41, Il2ra, Abl, Mpmv3, Rap1a-ps2, anx, Mpmv43, Ryr3, Rasl2-4, Adra2b, Avp, Glvr1, Il1a, Il1b, Mpmv28, Oxt, Pcsk2, a, Xmv10, Tcf4, Acra, Acra4, Ak1, Bdnf, bs, Cyct, Cyp24, Dbh, Fshb, Gcg, Gdf5, Gnas, Gpcr8, Grin1, Hcs4, Hior2, Hsp84-2, Idd12, Ilrn, Jund2, Kras3, Mc3r, Mpmv14, Mtv40, Mxil-rs1, Otf3-rs2, Ptgs1, Ptpra, Rapsn, Src, Svp1, Svp3, Tcf3b, Wt1, Xmmv71, Xmv48, Ccna, Fgf2, Fth-rs1, Csfm, Mov10, Egf, Acrb2, Cap1, Crh, Fim3, Fpsl1, Glut2, Gpcr2, Gria2, Hsd3b-1, Hsd3b-2, Hsd3b-3, Hsd3b-4, Hsp86-ps2, Idd3, Il2, Il7, Mpmv9, Mpmv20, Mtv4.8, Ngfb, Npra, Nras, Nras, Ntrk, Otf3-rs3, Otf3-rs4, Rap1a, Tshb, Xmmv22, Xmmv65, Mos, Rasl2-7, Lyr, Ifa, Ifb, Jun, azh, db, Ipp, Mpl, Do1, Ak2, Ccnb1-rs4, Cdc21l, Cga, Fgr, Foc1, Fpsl2, Gabrr1, Gabrr2, Gdf6, Glut1, Gnb1, Gpcr14, Grb2-ps, Grik3, Grik5, Hsp86-1ps4, Htr1da, Htr1db, Idd9, Ifa1, Ifa2, Ifa3, Ifa4, Ifa5, Ifa6, Ifa7, Ifa8, Ifa9, Ifa10, Lap18, Lmyc1, Mpmv19, Mpmv44, Mtv13, Mtv14, Mtv17, Nppb, Otf6, Otf7, Ril2, Ski, Tnfr2, Wnt4, Xmmv8, Xmmv23, Xmmv62, Xmv1, Xmv2, Xmv8, Xmv9, Xmv14, Xmv44, Xpa, Tec, Fgf5, Nos1, Tcf1, Epo, Gnb2, Flt1, Flt3, Ache, Adra2c, Adrbk2, Afp, Alb1, Ccnb1-rs1, Clock, Cyp3, Cyp3a11, Cyp3a13, Drd1b, Drd5, Fgfr3, Flk1, Gc, Gnrhr, Gpcr1, Hcs5, Hnfl, Htr5a, Il5r, Il6, Kit, Ltrm3, Mgsa, Mpmv7, Mpmv13, Mpmv23, Mtv32, Mtv41, Pdgfa, Pdgfra, Por, Txk, Xmmv3, Xmmv5, Xmmv52, Xmv17, Xmv28, Xmv34, Xmv38, Xmv45, Zp3, Trh, Raf1, Fth-rs2, Ntf3, Kras2, Pthlh, Mov1, Alox5, Braf2, Cfr, Egr4, Fpsl10, Fgf6, Gdf3, Ghrfr, Glut3, Grin2a, Hior3, Hoxa10, hop, Ica1, Il5r, Int41, Itpr1, Krag, Mad,

Met, Mi, Mtv8, Mtv23, Mtv29, Mtv33, Mtv34, Nkna, Npy , ob , Otf3-rs5, Tgfa, Tnfr1, Wnt2 , Wnt5B, Wnt7A, Xmmv27, Xmv24, Xmv61, Fosb, Ryr1, Ngfa, Ufo, Xrccl, Abpa, Abpga, Gabra4, Gas2, Acra7, Ccnb1-rs7, Egfbp3, Xmv30, Zp2, Fes, Pcsk3, Calc, Ccnb1-rs10, Pth, Ad, Bcl3, Cea, Cea2, Cea3, Cea4, Cea5, Cea6, Cebp, Dm9, Dm15, Drd4, Egfbp1, Egfbp2, Ercc2, Fgf3 , Fgfr2, Gabra5, Gabrb3, Gtx, Hcs1, Igflr, Igf2, Il4r, Ins2, Int40, Lhb, Mpmv1, Mtv1, Mtv35, Ngfg, Ntf5, Otf2, 2, Pkcc, Rasl4, Rras, Ryr, Svp2, Tcf3g, Tgfb1, tub , Xmmv31, Xmmv35, Xmmv73, Xmv33, Xmv53, Taz83, Adrb3, Junb, Jund1, Mel, Gpcr19-rs2, Agt, Cadp, Ccnb1-rs9, E, Fgfr1, Gas6, Gnb-rs1, Hcs2, Insr, Maf, Mov34, Mpmv21, Mpmv41, Mtv21, Mtnr1a, Plat, Rasl5-2, Rasl6, Sntb2, Xmmv29, Xmv12, Xmv26, Xmv62, Epdr, Gpcr13, Otf11, Pthr, Acra3, Acra5, Acrb4, Camkl, Cdc25Mm, Crbp, Crbp2, Csk, Cyp11a, Cyp19, Drd2, Ets1, Fli1, Gnai2, Gnat1, Gpcr6, Gria4, Hgfl, Hior1, Hpx, Hsp86-1ps3, Hst2, Idd2, Il1bc, Lag-rs1, Lap18-rs1, Mll, Mpmv27, Penk, Pgr , Rasl2-2, Tp11, Trf, Xmmv2 , Xmmv67, Xmv15, Xmv16, Xmv25, Xmv60, Mgf , Amh, Braf, Cdc2a, Dmdl, Estr, Fpsl3 , Fpsl4 , Fpsl5 , Gli, Gpcr17, Grik2 , Ifgr, Igfl, Mpmv5, Mpmv12, Mpmv40, Myb, Oprm, Pg, Pmch, Ros1, Xmv31, Xmv51, Xmv54, Camk2b, Egfr , Int6, Lif, Mtv44, Ews, Csfgm, Flt4, Il3, Il4, Il5, Irf1, Gria1, Glut4, Crhr, Csfg, Mov9, Xmv20, Acrb, Mpmv4, Mpmv15, Ngfr, Nos2, Rara, Taz4, Tcf2, Xmv42, Mtv3, Adra1, Crko, df, Erbb2, Gabra1, Gabra6, Gabrg2, Gh, Glra1, Grb2, Hnflb, Hsp86-ps1, Idd4, Igfbp1, Igfbp3, Il13, Int4, Mpmv2, Mpmv8, Mpmv18, Mtv45, nu, Pkca, Rab1, Rel, Shbg, Tcf7, Thra, Tnz1, Trp53, Wnt3, Wnt3A, Xmv4, Xmv5, Xmv47, Xmv49, Xmv63, Akt, Amh-rs4, Ccs1, Fpsl6, Fos, Gdf7, Hcs3, Hsp70-2, Hsp84-3, Hsp86-1, hyt, Ltrm1, Max, Mpmv11, Mpmv24, Mtv9, Mtv30, Pomcl, Tcf3a, Tda2, Tgfb3, Tpo, Tshr, Xmmv21, Xmmv25, Xmmv34, Xmmv50, Gli3, Xmv55, Ryr2, Inhba, Gas1, Pcsk1, Amh-rs2, Ccnb1-rs6, Ccnb1-rs13, Crhpb, Dat1, Drd1a, Fgfr4, Fpsl7, Fim1, Gpcr15, Gpcr18, Hbvi, Hilda, Htr1a, Idd11, Il9, Ltrm4, Mak, mes, Pl1, Pl2, Prl, Ral, Rasa, Srd5a1, Tpbp, Xmv13, Xmv27, Rarb, Rbp3, Htr2, Rb1, Acra2, Camkg, Cchl1a2, Ccnb1-rs5, Ccnb1-rs12, Gnrh, Mtv11, Nras-ps, Otf3-rs6, Plau, Ptprg, Trp53-ps, Wnt5A, Xmv19, Ghr, Il7r, Lifr, Mlvi2, Prr , Myc , Ril1, cog, Amh-rs7, Il2rb, Pdgb, Acr, CP2, Rarg, Sp1-1, Wnt1 , Afr1, Atf4, Bzrp, Ccnb1-rs11, Cyp11b, Il3rb1, Il3rb2, Ins3, Itga, Mlvi1, Mlvi3, Mtv36, Pdggfec, Svp5, Tef, Trhr, Wnt7B, Xmmv55, Xmmv72, Xmv37, Tnp2, Ets2, Casr, Chuck-rs1, din, Drd3, Erg, G22p1, Gap43, Gas4, Grik1, Htr1f,

Ifgt, Int53, Ltrm2, Mpmv17, Mtv6, Mtvr1, Pit1, Xmv3, Xmv35, Xmv50, Igf2r, Mas, Tcd3, Glp1r, Idd1, Tla, Aeg1, Ccnb1-rs3, Cdc2b, Csi, Cyp21, Cyp21-ps1, Fpsl8, Gna-rs1, Gpcr19-rs1, Grr1, Grr2, Hom1, Hsc70t, Hsp70, Hsp70-1, Hsp70-3, Hsp84-1, Hst1, Hst4, Hst5, Hst6, Hye, Int3, Itpr3, Lap18-rs2, Otf3, Ptprs, Rab11b, Rasl2-1, Rasl2-3, Rasl3, Rrs, Rxrb, Tas, Tcd1, Tcd2, Teral, Tla-rs, Tnfa, Tnfb, Tpx1, Tpx2, Xmmv15, Xmv36, Xmv57, Csfmr, Pdgrfb, Adrb2, Apc, Camk2a, Camk4, Dcc, Fgf1, Gnal, Gpcr7, Grl1, Grp, Hsp74, Mcc, Mtv2, Mtv38, Ptpn2, Tpl2, Xmv22, Xmv23, Xmv29, Fth, Csfgmra, Mxi1, Adra2a, Adrb1, Adrbk1, Chuck, Cyp17, Gna14, Gnb-ps1, Hcs6, Htr7, Ide, Ins1, Lpc1, Pomc2, Seao, Tlx1, Xmmv42, Xmv18, Tcfe3, Araf, Avpr2, mdx, Ar, Zfx, Otf9, Ccg1, Ccnb1-rs8, Fpsl9, Gabra3, Glra2, Glra4, Gria3, Grpr, Hsp74-ps1, Hst3, Htr1c, Il2rg, Mov14, Mov15, Mtv28, Otf3-rs8, Sts, Sxa, Sxr, Xta, Tdy, Hya, Zfy1, Zfy2, Mov15, Mov24, Mtv31, Mtv42, Sdma, Spy, Sts, Sxa, Sxr, XmmvY, Xmv7, Xmv11, and Xmv40.

[0067] Examples of *Phaseolus vulgaris* genes include: Acc, ace, Adk, Am, Amv-1, Amv-2, Ane, aph, Arc, Are, arg, Arl (Arc), asp, B, bc-u, bc-1<sup>1</sup>, bc-1<sup>2</sup>, bc-2<sup>1</sup>, bc-2<sup>2</sup>, bc-3, Bcm, Beg, Bip, blu, Bpm, Bsm, By-1, By-2, C, C/c, c<sup>cr</sup>, C<sup>cir</sup>, C<sup>ma</sup> (M, R<sup>ma</sup>), C<sup>r</sup>, C<sup>res</sup>, C<sup>rho</sup>, C<sup>st</sup>, [C<sup>st</sup> R Acc] (Aeq), c<sup>u</sup> (inh, i<sub>e</sub>), [c<sup>u</sup> Prp<sup>i</sup>] (Prp, c<sup>ui</sup>, Nud), [c<sup>u</sup> prp<sup>st</sup>] (prp<sup>st</sup>), [C Prp] (Prp), c<sup>v</sup>, [C R] (R), [C r] (r), Ca, Cam, Cav, cc, chl, cl, cml, Co-1 (A), Co-2 (Are), Co-3 (Mexique 1), Co-3<sup>2</sup>, Co-4 (Mexique 2), Co-5 (Mexique 3), Co-6, Co-7, cr-1 cr-2, cry, cs, Ct, ctv-1 ctv-2, cyv (by-3), D (Can, Ins), Da, Db, def, dgs (gl, le), dia, Diap-1, Diap-2, diff, dis, Dl-1 Dl-2 (DL<sub>1</sub> DL<sub>2</sub>), do, ds (te), dt-1<sup>a</sup> dt-2<sup>a</sup>, dt-1<sup>b</sup> dt-2<sup>b</sup>, dw-1 dw-2, Ea Eb, ers (restr), ers-2, Est-1, Est-2, exp, F, Fa, fast, Fb Fc, fa fb fc, Fcr, Fcr-2, fd, Fe-1 Fe-2, Fin (in), Fop-1, Fop-2, Fr, Fr-2, G (Flav, Ca, Och), Ga, gas, glb, Gpi-c1, Gr, Hbl (L<sub>HB-1</sub>), Hbnc (SC<sub>HB-1</sub>), Hbp (PD<sub>HB-1</sub>), hmb, Hss, Hsw, Ht-1 Ht-2 (L-1 L-2), I, Ia Ib, ian-1 ian-2 (ia), lbd, ico, Igr (Ih), ilo, ip, iter, iv, iw, J (Sh), Ke, L, la, Lan, Ld, Lds (Ds), Lec, Li (L), lo, lr-1 lr-2, mar, Me, Mel (Me), Mel-2 (Me-2), mel-3 (me-3), Mf, mi, mia, Mic (Mip), miv, Mrf, Mrf<sup>2</sup>, mrf, ms-1, Mue, mu mutator, Nag, Nd-1 Nd-2 (D-1 D-2), nie, nnd (sym-1), nnd-2, No, nts (nod), Nudus, ol, P, p<sup>gri</sup> (Gri, v<sup>pal</sup>), pa, pc, pg (pa), Pha, Pmv, ppd (neu), Pr, prc (pc), Prx, punc, ram, Rbcs (rbcS), rf-1, rf-2, rf-3, rfi (i), Rfs (m), Rk, rk, rk<sup>d</sup> (lin), rn-1 rn-2 (r r), rnd, Ro, Sal, sb, sb<sup>ms</sup>, sb-2, sb-3, sil, Skdh, sl, Smv, St, Sur, sw-1 sw-2, T, t (z-1), Th-1 Th-2, Tm, To, Tor (T), Tr, tri, trv, Ts, tw, uni, Uni-2, uni<sup>nde</sup>, uni<sup>nie</sup>, Ur-1, Ur-2, Ur-2<sup>2</sup>, Ur-3 (Ur-3, Ur-4), Ur-3<sup>2</sup>, Ur-4 (Up-2, Ur-C), Ur-5 (B-190),

Ur-6 (Ur<sub>a</sub>, Ur-G), Ur-7 (R<sub>B11</sub>), Ur-8 (Up-1), Ur-9 (Ur<sub>p</sub>), us, V (Bl), v<sup>lac</sup> (Cor), v, var, vi (vir<sub>l</sub>), wb, Wmv, X<sup>su</sup>, y, and Z.

[0068] Examples of *Saccharomyces cerevisiae* genes include: PRE3, PUP1, PUP3, PRE2, PRE10, PRE1, PRE8, SCL1, PUP2, PRE5, PRE7, PRE4, RPT2, RPT3, RPN3, RPN11, RPN12, RPT6, RPN1, RPN2, RPT1, RPT5, RPT4, SKI6, RRP4, DIS3, TSC10, RAT1, GND1, EXO70, ERG10, ACC1, RPP0, ACT1, ARP100, ARP3, PAN1, ARP2, ARP4, ARP9, SPE2, CYR1, ALA1, TPS1, TUB1, ABF1, DED81, NIP1, YHC1, SNU71, ATM1, MAK5, ROK1, DED1, SPB4, AUR1, PSE1, ALG1, TUB2, BPL1, MSL5, ERG24, ERG26, ERG25, CMD1, HCA4, SHE9, SHE10, CAK1, PIS1, CHO1, CDS1, ESR1, NUD1, CDC47, CDC13, CDC37, CDC1, CDC4, CDC20, CDC6, CDC46, CDC3, KAR1, BBP1, HRP1, CCT2, CCT3, HSP10, SMC1, SMC2, CHC1, CFT2, CLP1, COP1, SEC26, SEC27, RET2, SEC21, COF1, CCT4, CCT1, CCT6, SEC24, SEC7, PCF11, RNA15, RNA14, FIP1, YSH1, TFB4, TSM1, APC2, APC5, SEC31, TAF47, TAP42, MPP10, CDC53, CKS1, CDC28, KIN28, CNS1, ERG11, DBP10, DBP8, PRO3, DYS1, ALR1, TID3, DNA2, SSL2, RAD3, RFA3, RFA2, RFA1, RFC4, RFC5, RFC3, RFC2, RFC1, TOP2, RAP1, RPC25, PRI2, PRI1, POL1, POL12, HUS2, CDC2, POL2, DPB2, RPB10, RPA135, RPA190, RPA43, RPB8, RPO26, RPB5, RPC40, RPC19, SRB7, SRB4, RGR1, RPB11, SRB6, RPB2, RPB7, RPO21, RET1, RPO31, RPC31, RPC34, RPC53, RPC82, RPB12, RPB3, DPM1, DIP2, RNT1, CDC8, CDC14, DUT1, UBA2, UBA1, UBC9, CDC34, ENP1, ERD2, SSS1, SEC61, SEC63, SEC62, GNA1, GPI8, DAM1, DUO1, IRR1, PRP3, TIM9, HSH49, SUP35, EXM2, MEX67, ERG9, ERG20, FAS2, FAS1, NOP1, FAD1, AOS1, FBA1, NCB2, BRN1, TUB4, GDI1, GOG5, SRM1, CDC25, SPT16, YIF2, BET4, CDC43, MRS6, BET2, PRO1, GLN1, GLN4, GRS1, YIP1, FOL2, GPA1, CDC42, SAR1, YPT1, SEC4, GSP1, TEM1, RHO1, CDC24, RNA1, GUK1, VMA16, PMA1, HKR1, SIS1, MGE1, HSP60, HSF1, HAS1, MOT3, HTS1, ESA1, HSL7, HOM6, RIB7, SLY1, CSL4, PUR5, CSE1, IPP1, MDM1, USO1, SOF1, MAK11, LAS1, TEL2, DPB11, SGD1, FAL1, MTR3, MTR4, SPP2, SIK1, RRP7, POP4, RRP1, POP3, BFR2, CDC5, NRD1, MET30, MCM6, RRP46, SAS10, SCC2, ECO1, PRP43, BET3, BET5, STN1, NFS1, IDI1, SRP1, KAP95, CBF2, SKP1, CEP3, CTF13, ERG7, KRS1, PSA1, PMI40, ALG2, SSF1, MED7, RSC4, CDC54, MCM2, AFG2, ERG12, MVD1, CDC48, MHP1, ERV1, SSC1, TIM44, TIM17, TIM23, TOM22,



TOM40, MAS1, MCD1, MMC1, STU1, JAC1, ABD1, CEG1, PAB1, MTR2, SEC16, ROT1, INO1, MLC1, MYO2, GPI2, SPT14, NAT2, NMT1, TRM1, NCP1, NBP1, ACF2, SPP41, NUT2, LCP5, PRP19, NMD3, RFT1, NNF1, NDC1, CRM1, KAR2, NIP29, NAB2, NIC96, NUP145, NUP49, NUP57, NUP159, NSP1, NUP82, CDC39, NPL4, POP7, NTF2, MAK16, NPL3, NOP2, NOP4, NHP2, NOP10, GAR1, NBP35, WBP1, STT3, SWP1, OST2, OST1, ORC1, ORC6, ORC5, ORC4, ORC3, RRR1, SAT2, PWP2, PEX3, TOR2, PIK1, SEC14, STT4, MSS4, PCM1, GPM1, SEC53, ERG8, YPD1, PAP1, NAB3, RRN7, SEN1, CFT1, PRP11, PRP21, PRP39, PRP24, PRP9, SLU7, PRP28, PRP31, IFH1, PTA1, SUB2, FMI1, MAS2, ESS1, PFY1, POL30, POP1, PDI1, RAM2, CDC7, SMP3, CDC15, YTH1, QRI2, YAE1, SFI1, SEC1, BET1, SEC6, SEC13, SEC2, SEC8, CBF5, CDC19, , YRB1, RHC18, DBF4, SDS22, MCM3, CEF1, ALG11, GAA1, MOB1, NIP7, TIP20, SEC5, SEC10, GPI10, RRP3, CDC45, DIB1, MIF2, HOP2, PBN1, NOP5, RPP1, POP5, POP8, POP6, ERO1, MPT1, DNA43, ESP1, SMC3, LST8, STS1, RPM2, RNR1, RNR2, RNR4, RPS20, RPL25, RPL3, RPL30, RPL32, RPL37A, RPL43A, RPL5, RPL10, RPS3, CET1, YRA1, SNM1, GLE1, DBP5, DRS1, DBP6, BRR2, RRN3, RRN6, RRN11, MED6, PRP16, RPR2, DIM1, RRP43, RRP42, RRP45, SEC20, BOS1, CDC12, GLC7, PKC1, IPL1, SGV1, NRK1, RAD53, LCB2, LCB1, MPS1, SES1, SPC3, SEC11, RIO1, ARP7, NEO1, YJU2, POB3, ARH1, IQG1, HRT1, HYM1, MAK21, FUN20, FUN9, NBN1, STB5, YIF1, SMX4, YKT6, SFT1, SMD1, PRP6, LSM2, NUF1, SPC97, SPC42, SPC98, CDC31, SPC19, SPC25, SPC34, SPC24, NUF2, PRP40, MCD4, ERG1, SMC4, CSE4, KRR1, SME1, TRA1, RLP7, SCH9, SMD3, SNP2, SSF2, SPC72, CDC27, CDC23, CDC16, APC1, APC11, APC4, ARC19, RPN6, RPN5, RSC6, RSC8, STH1, SFH1, TIM12, TIM22, TIM10, SQT1, SLS1, JSN1, STU2, SCD5, SSU72, ASM4, SED5, UFE1, SYF1, SYF2, CCT5, TBF1, TOA2, TOA1, SUA7, TAF90, TAF61, TAF25, TAF60, TAF17, TAF145, TAF19, TAF40, TAF67, TFA2, TFA1, FCP1, TFG1, TFG2, TFB1, CCL1, SSL1, TFB3, TFB2, PZF1, BRF1, TFC5, TFC4, TFC3, TFC7, TFC6, TFC1, SPT15, THI80, THS1, SPT6, SPT5, ROX3, REB1, MCM1, MED4, MOT1, MED8, EFB1, YEF3, SUI1, CDC95, TIF11, SUI3, GCD11, SUI2, GCD6, GCD7, GCD2, GCD1, RPG1, GCD10, PRT1, TIF34, CDC33, TIF5, SUP45, GCD14, TIM54, SEC17, TPT1, TRL1, CCA1, SEN54, SEN2, SEN15, SEN34, WRS1,

**SECRET**

[0069] The vaccinia virus of the invention can be used in a method of treatment of man or animal body. Such treatment includes a method of treating the growth of stic cells which comprises administering to a patient in need of treatment the vaccinia of the invention. The recombinant vaccinia virus of the invention may or may not an exogenous nucleic acid sequence.

[0070] For use of the expression vectors in therapy, the expression vectors will be packaged into viral particles and the particles delivered to the site of the tumor. Particles may be delivered to the tumor by any suitable means at the disposal of the artisan.

[0071] Recombinant infectious vaccinia viruses for use in man can be prepared as described in Br. Med. Bull. 25: 131-135 (1969). Preparations suitable for vaccination often contain  $10^6$  to  $10^8$  plaque forming units per 0.05 ml.

[0072] One suitable route of administration is by injection of the particles in a solution. While it is possible for the particles to be administered alone it is preferable to present them as pharmaceutical formulations. The formulations comprise a particle, or with one or more acceptable carriers thereof and optionally other therapeutic agents. The carrier or carriers must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipients thereof.

[0073] Particles may also be administered by regional perfusion or direct arterial direction, or direct injection into a body cavity (intracavitary administration), for example by intra-peritoneum injection.

[0074] It is also known that muscle cells can take up naked DNA and thus may be treated using an expression vector of the invention in which naked DNA is injected into the sarcoma.

**0075]** Formulations suitable for parenteral or intramuscular administration aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, bacteriostats, bactericidal antibiotics and solutes which render the formulation with the blood of the intended recipient; and aqueous and non-aqueous sterile

09091724.14304

suspensions which may include suspending agents and thickening agents and preserving agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example water, for injections, immediately prior to use.

[0076] It should be understood that in addition to the ingredients particularly mentioned above the formulations may include other agents conventional in the art having regard to the type of formulation in question. Of the possible formulations, sterile pyrogen-free aqueous and non-aqueous solutions are preferred.

[0077] The doses may be administered sequentially, eg. at daily, weekly or monthly intervals, or in response to a specific need of the patient. Preferred routes of administration are parenteral injection, typically intradermal injection or intramuscular injection or intratumoral injection.

[0078] Typically, the particle will be administered to the patient and then the uptake of the vector by the infected cells monitored, for example by recovery and analysis of a biopsy sample of targeted tissue.

[0079] It will be appreciated that the actual preferred amounts of active compound in a specific case will vary according to the specific compound being utilized, the particular compositions formulated, the mode of application, and the particular situs and organism being treated. Dosages for a given host can be determined using conventional considerations, e.g., by customary comparison of the differential activities of the subject compounds and of a known agent, e.g., by means of an appropriate, conventional pharmacological protocol.

[0080] Chemotherapy will normally involve parenteral administration. Administration by the intravenous route is frequently found to be the most practical. For glioblastoma the route is often intratumoral.

[0081] Tumors which may be treated using the expression vector of the present invention include any tumors capable of being treated by the expression vector of the invention and thus are not limited to any one particular class of tumors. Particularly suitable

[illegible]

**090917Z** **FM** **USCINCPAC** **TO** **CINCPACFLT**

**090917Z**

**090917Z**

**090917Z** **FM** **USCINCPAC** **TO** **CINCPACFLT**

**090917Z** **FM** **USCINCPAC** **TO** **CINCPACFLT**

[illegible][illegible]

**090917** **Z** **E** **I** **J** **B**

[0086] The pSC16 plasmid was cleaved with *AccI* and then subjected to bidirectional BAL 31 digestion and Klenow enzyme repair. The DNA was modified by the addition of *BglII* linkers, circularized with T4 DNA polynucleotide ligase, and used to transform competent HB101 bacteria. Minipreparations of plasmids from isolated colonies were characterized by digestion with *BglII* and *HindIII*. A plasmid, pSC18, was identified in which the *AccI* site had been destroyed by a deletion of approximately 250 base pairs and a *BglII* site had been inserted. This plasmid was treated sequentially with *BglII* and calf intestinal phosphatase and ligated to an appropriately modified *XbaI-SmaI* fragment from pSC10, described in Chakrabarti, et al., Mol. Cell. Biol. 5:3403-3409 (1985), which contained the vaccinia virus 11K promoter coupled to the *E. coli lacZ* gene ( $\beta$ -galactosidase cassette). Competent JM109 cells were transformed, and blue colonies ( $\beta$ -Gal+) were isolated. One colony yielded a plasmid (pSC20) which, on characterization with *HindIII* and *ClaI* restriction endonucleases, showed a pattern consistent with the  $\beta$ -galactosidase cassette having been inserted into the disrupted coding region of the VGF gene and in the same transcriptional orientation.

[0087] The pSC20 plasmid was then used to introduce the mutated VGF gene into the vaccinia virus genome by homologous recombination, resulting in the production of a VGF- phenotype vaccinia virus. To accomplish this a 25-cm<sup>2</sup> flask of CV-1 cells (monkey kidney cells) were grown to confluency in a 6-well plate. The CV-1 cells were infected with 0.05 plaque forming units (pfu) per cell of wild-type vaccinia virus. These cells were then transfected with 5-10  $\mu$ g (in <50  $\mu$ l) of plasmid pSC20 using the calcium phosphate-precipitation method. After a 30 minute incubation at room temperature, 9 ml of complete MEM-10 medium was added and the cells were then incubated for 3 to 4 hours at 37°C. Following this incubation period, the medium was removed and replaced with 5 ml complete MEM-10, and then the cells were incubated for 2 days at 37°C. Following the incubation the cells were dislodged with a scraper and transferred to a 15-ml conical centrifuge tube. The tube and its contents were centrifuged for 5 minutes at 1800 x g (2500 rpm in a Sorvall H-6000A rotor), at 5° to 10°C. The medium was then aspirated and discarded. The cells were then resuspended in 0.5 ml complete MEM-2.5. The cell suspension was lysed by performing three freeze-thaw cycles, each time by freezing in a dry ice/ethanol bath, thawing

in a 37°C water bath, and vortexing. The resulting lysates were then stored at -70°C until needed for selection of the VGF- virus.

[0088] To isolate VGF- phenotype virus, CV-1 cells were cultured to a confluent monolayer and then trypsinized and resuspended in culture medium. The cells were cultured and placed in a 6-well tissue culture plate at  $5 \times 10^5$  cells/well in a final volume of 2 ml/well. The plate was incubated until the cells were confluent (<24 hours). Following establishment of confluency, 100  $\mu$ l of transfected cell lysate and 100  $\mu$ l of 0.25 mg/ml trypsin was mixed and vortexed vigorously. The mixture was incubated for 30 minutes at 37°C with periodic vortexing at 5 to 10 minute intervals. Following incubation the mixture was vortexed for 30 seconds on ice. Four 10-fold serial dilutions (ranging from  $10^{-1}$  to  $10^{-4}$ ) were made for use in the assay. The medium was then aspirated from the cell monolayers and infected with 1.0 ml diluted lysate per well. This solution was incubated for 2 hours with rocking at 30 minute intervals.

[0089] Before the 2 hour infection was finished, 2% LMP agarose (Life Technologies; Rockville, MD) in water (1.5 ml times the number of wells) was melted and placed in a 45°C water bath to cool. Plaque medium was prepared by making 2X culture medium and warming it to 45°C. A first overlay of agar was then prepared by mixing an equal volume of 2% LMP agarose and plaque medium. The viral inoculum was then removed and an overlay of 3 ml per well was placed in each well and allowed to solidify at room temperature or 4°C for 2 days.

[0090] A second agarose overlay was then prepared by mixing equal volumes of 2% LMP agarose (1 ml x number of wells, melted and cooled to 45°C as described above) and 2X plaque colorimetric medium (1 ml times the number of wells, warmed to 45°C) with 1/100 volume of 10 mg/ml neutral red. An additional 1/120 volume of 4% Xgal was added to the plaque colorimetric medium. Each well was then overlayed with 2 ml of this second agarose solution, allowed to solidify, and incubated overnight to permit color development. Plaques with blue coloring indicated the presence of an active  $\beta$ -galactosidase gene and thus an interrupted VGF gene.

[0091] Blue plaques were isolated by taking an agar plug of a blue plaque and placing it into a tube with 0.5 ml of culture medium. Each tube was vortexed and subjected

to three rounds of freeze-thawing as described above. At the end of the third round the tube and its contents were sonicated in ice-water for 20-30 seconds. The resulting contents of the tube were used to perform successive rounds of plaque purification. These subsequent rounds of purification are necessary in view of the vaccinia virus genome's ability to transfer the genotype of one VGF gene to the other copy within the vaccinia virus genome. Because of this characteristic of vaccinia virus, it is important to select a mutant vaccinia virus clone where both VGF genes are mutated, rather than only one. Often by the fourth round of plaque purification all of the plaques were blue, suggesting that the  $\beta$ -galactosidase cassette was present in both inverted terminal repeats of the vaccinia virus genome. One plaque was designated VSC20. This plaque was used in the construction of the VGF-/TK- double mutant virus.

[0092] Once the VSC20 plaque was selected it was amplified to create a virus stock. To amplify the virus from the selected plaque, the plaque plug was selected, resuspended and sonicated as described above. A confluent monolayer of CV-1 cells was then infected with the solubilized plaque material and incubated for 2 days. The infected culture cells were then scraped, lysed, and centrifuged to remove the particulate matter. This material was then used in subsequent scale up procedures to produce a substantial virus stock for storage.

#### TK-/VGF- Double Mutant Vaccinia Virus

[0093] The vaccinia virus VSC20 described above was used in the generation of the VGF-/TK- phenotype vaccinia virus using standard techniques well known in the art. For example, such techniques are described in detail in Moss & Earl, Current Techniques in Molecular Biology, (Ed. Ausubel, et al.) Unit 16.15-16.19 (1998). Purified VSC20 virus (VGF- phenotype) was used to infect HuTK- 143B cells grown to confluency in a 6-well plate. The cells were infected with  $6 \times 10^6$  plaque forming units (pfu) of VSC20 and incubated for 2 hours in the presence of 5-bromodeoxyuridine (BrdU), a nucleotide analog that incorporates into DNA in the presence of a functional TK gene, and results in cell death.

[0094] When viral cytopathic effects were observed in the cultured cells (rounding up of cell layer), the culture media was aspirated and the cells were resuspended in complete DMEM with 2.5 % serum. Dilutions of cells ( $10^{-1}$  to  $10^{-3}$ ) were used to infect

confluent HuTK- cells in presence of BrdU in a plaque assay as described above. Only TK negative phenotype viruses can grow in the presence of BrdU. Viral plaques that developed were thus TK negative. Plaques that appeared in this assay were then picked and used to infect further HuTK- cells for 3 cycles. Multiple plaques were then expanded on CV-1 cells.

[0095] DNA was extracted from these cells using methods well known in the art. The plaque DNA isolated was used in various molecular biology techniques to determine the presence or absence of VGF and TK mutations. For example the polymerase chain reaction (PCR) reaction was performed to examine the size of the TK and VGF genes. The reaction was performed using standard techniques well known in the art. Briefly, 1 µl of isolated DNA and 1µl of each primer (TK sense primer: 5'-GATCTTCTATCTCGGTTTCCTCAC-3' (SEQ ID NO. 1); TK antisense primer: 5'-GATCGATAATAGATACGGAACGGG-3' (SEQ ID NO. 2); VGF sense primer: 5'-CTGATGTTGTTGTTTCGTCGC-3' (SEQ ID NO. 3), VGF antisense primer: 5'-GGTAGTTTAGTTCGTCGAGTGAACC-3' (SEQ ID NO. 4))) were added to 47 µl of PCR Supermix (Gibco BRL, Gaithersburg, MD). 25 cycles, each consisting of 15s denaturing (94 C), 30s annealing (55 C) and 2 minute extension (72 C) were performed. The results of the PCR showed the absence of the VGF genes and a normal sized TK gene. Subsequence analysis (Automated sequencing, ABI PRISM DNA Sequencing Kit, Perkin Elmer Applied Biosystems; Foster City, CA) of the TK gene revealed a C to A base change resulting in an alanine to glutamic acid amino acid change. The resultant amino acid alteration produced a functionally inactive TK protein, and thus a TK- phenotype. The VGF gene remained disrupted. The isolated double-deleted vaccinia virus (VVDD) was expanded in HeLa cells to 10<sup>9</sup> pfu per ml and used in subsequent experiments.

#### Example 2

##### An Alternative Method of Creating a Double Mutant Vaccinia Virus

[0096] In another embodiment, the double mutant vaccinia virus is generated by introducing both VGF and TK mutations through homologous recombination. The VSC20 virus discussed in Example 1 is used to infect HuTK- 143B cells as described above. These cells are contacted with a vaccinia virus shuttle vector containing a mutated TK gene.



TEFTT "T24T5550

[0097] The pSC65 vector is used as the source of the TK gene and is described in Chakrabarti, et al., (1997). The *Xba*I site of the pSC65 vector within the flanking region of the thymidine kinase gene is abolished following digestion using the Klenow enzyme and dNTP filling. A segment of the plasmid is amplified using the PCR reaction to introduce a *BSS*II site between the TK flanking region, allowing for the insertion of the multiple cloning site (MCS) of pBluescript KS II (+) (Stratagene; La Jolla, CA). To prevent possible antisense transcripts driven by the native vaccinia thymidine kinase promoter, two primers are designed to encode an early termination signal (TTTTTNT) for all three possible open reading frames (5'-ATC GGA GCT CTT TTT ATC TGC GCG GTT AAC CGC CTT TTT ATC CAT (SEQ ID NO. 5) and 5'-ATC GTC TAG ACT CCA CAA TAA AAA CAG ATC ACC TGA TGG ATA AAA (SEQ ID NO. 6)). The primers are hybridized, and dNTPs and Klenow enzyme are added to the reaction mixture. After restriction with *Sac*I/*Xba*I, the termination signal is gel isolated and ligated into the above plasmid. Gpt is cut out of pBSgpt Puhlmann, et al., Cancer Gene Therapy, (In press), using *Xho*I and *Eco*RV and inserted into the MCS of the resulting plasmid pCB022. A promoter cassette consisting of a vaccinia virus p7.5 early/late and a synthetic early/late promoter, as described in Chakrabarti, et al., BioTechniques 23:1094-1097 (1997), in sense and antisense direction is inserted immediately upstream so that the gpt- gene is under control of the p7.5 promoter. A gene of interest, for example the luciferase gene from pGEM-luc (Promega; Madison, WI), is inserted into pCB022 to be driven by the vaccinia synthetic early/late promoter.

[0098] The shuttle vector is introduced into a vaccinia virus host cell in conjunction with the VSC20 virus with the VGF- phenotype as described above. Virus progeny are screened for TK- phenotype using BrdU selection described in Example 1. As a result of homologous recombination, a TK-, VGF- negative phenotype is obtained.

### Example 3

#### An In Vitro Comparison of the Double Mutant Vaccinia,

#### TK-, VGF-, and Wild-type Viruses

[0099] To compare the growth characteristics of the double mutant VGF-/TK- vaccinia virus with those of the single TK- and VGF- mutants and the wild-type virus, each were grown in confluent and non-confluent cells. Six well plates of confluent and non-

confluent (dividing) NIH3T3 (growth factor dependent) were infected with 167 pfu per well of one of the following vaccinia viruses: F13 (wild-type)(Blasco & Moss, *Gene* 158:157-162 (1995)), VJS6 (thymidine kinase deleted)(Carroll, et al., *Vaccine* 15:387-394 (1997)), VSC20 (vaccinia growth factor deleted), or VVDD (double deleted). At 24, 36 and 48 hours the growth media was removed, and the cells were harvested as described above. The total viral titers was measured in a plaque assay described below.

[0100] The plaque assay was performed as described above with certain modifications. The cells were resuspended in 2 ml of DMEM-2.5 and subjected to one freeze-thaw cycle. Five hundred microliters (500  $\mu$ l) of cell lysate was used to infect CV-1 cells, as discussed above. After 1 hour an overlay (1:1 1.5% agar to DMEM-10) was placed over a confluent layer of CV-1 cells. Three (3) days later cells were stained with crystal violet and plaques counted.

[0101] Shown in FIGURE 1 are the plaque assay results (pfu measurements made in CV-1 cells) for each virus at the three time points for growth in NIH3T3 cells. In FIGURE 1A (Non-Confluent Cells), the wild-type form of the virus showed the highest level of viral replication. The VGF- and TK- single mutants and the double mutant forms of the virus all replicated in the dividing cells. In FIGURE 1B (Confluent Cells) the wild-type form of the virus produced the highest levels of viral progeny in these non-dividing cells, however, the VGF-, TK-, and double mutants failed to show statistically significant replication in these non-dividing cells. Accordingly, the mutations of the present invention restrict viral replication to dividing cells.

#### Example 4

##### An In Vivo Comparison of the Double Mutant Vaccinia, TK-, VGF-, and Wild-type Viruses

[0102] To investigate the effect of the double mutations and the corresponding VGF-/TK- phenotype of the mutant vaccinia virus on pathogenesis, female nude mice were injected with  $10^5$  MC38 cells (murine colon adenocarcinoma cells) subcutaneously. It is thought that vaccinia virus replicates in MC38 tumor cells and will eventually cause the death of the immunocompromised mouse. On day 10, when tumors were approximately 100  $\text{mm}^3$  in volume, the mice received either  $10^7$  or  $10^8$  pfu of one of the four viruses discussed in

Example 3. Specifically, F13 (wild-type), VJS6 (thymidine kinase deleted), VSC20 (vaccinia growth factor deleted), or VVDD (double deleted) were used in this experiment and injected intraperitoneally. The mice were observed for survival after viral infection.

[0103] As shown in FIGURES 2 and 3, there are substantial differences in survival between mice challenged with F13, those challenged with the two single mutation forms of the virus, or the double mutant vaccinia virus (VVDD). In FIGURE 2, mice that were challenged with the four virus types showed a marked difference relative to the number of mutations present in the viral genome. Comparing the point at which each of the four virus infected groups had 50% of their members still living, the wild-type virus had killed half of the mice treated with this form of the virus by approximately day 6. The VSC20 and VJ56 virus infected mice reached the 50% survival point on about days 26 and 27 (respectively). Mice infected with the double mutant virus showed significantly longer survival times and did not reach the 50% survival point until about day 54. These results indicate that mice harboring tumor cells survived viral infection for significantly longer periods of time when infected with the double negative phenotype form of the vaccinia virus.

[0104] Mice challenged with tumor cells as described above and exposed to VVDD substantiate the reduced toxicity of the vector. For example, half of the mice that were infected with the wild-type form of the virus were dead by day 6, and after that time point all of the remaining mice died. In contrast, mice exposed to cancer cells and to the double mutant vaccinia virus showed a substantial difference in survival relative to those mice infected with the wild-type form of the virus. As shown in FIGURE 3, 50% of the mice infected with the double negative phenotype mutation and the cancer cells survived until day 58. The mice infected with the TK- and VGF- single mutation viruses reached the 50% survival point on days 35 and 21, respectively. These results further confirm the observation that mice harboring dividing cancer cells survive longer when infected with the double mutant vaccinia virus of the present invention.

### Example 5

#### An *In Vivo* Comparison of the Wild-Type, Single Mutant, and Double Mutant Vaccinia

##### Expressing the Green Fluorescent Protein

[0105] This Example discusses an examination of host survival after administration of various forms of vaccinia. To make this comparison, less than  $10^7$  pfu of F13 (wild type), VJS6 (TK-deleted), VSC20 (VGF-deleted), and VVDDEGFP (TK- and VGF-deleted expressing green fluorescent protein) were injected intraperitoneally (in 2ml of HBSS/0.1%BSA, Calbiochem, La Jolla, CA) into six week old female, athymic (C57Bl/6) mice obtained from the National Institutes of Health small animal facility (Frederick, MD). The animals were examined daily with survival as an end point. Animals who became moribund due to viral pathogenicity were euthanized. The percent survival is graphed as a function of time. The data from this experiment is shown in FIGURE 4.

[0106] As shown in FIGURE 4, 50% of the mice infected with the double negative phenotype mutation and the cancer cells survived until day 141. The mice infected with the TK- and VGF- single mutation viruses reached the 50% survival point on days 38 and 19, respectively. These results further confirm the observation that mice survive longer when infected with the double mutant vaccinia virus of the present invention, even when that mutant virus is expressing an exogenous nucleic acid sequence.

### Example 6

#### An *In Vivo* Comparison of the Neurovirulence of Double Mutant Vaccinia, TK<sup>-</sup>, VGF<sup>-</sup>, and Wild-type Viruses in Brain Tissue

[0107] A comparison of neurovirulence, or ability to replicate in neurological tissues, in mice was performed using the viruses discussed in Example 5. Median viral titers were measured after  $10^7$  pfu of the viruses were administered IP to tumor bearing host animals. Virus titers were measured as a function of pfu/mg protein from brain and tumor tissue harvested from infected host animals. The results are shown in FIGURE 5.

[0108] As shown, the double mutant vaccinia virus VVDDEGFP failed to replicate appreciably in brain tissue. In contrast, the wild-type and single mutant forms of vaccinia were able to replicate in the brain tissue of the host animals. All forms of the virus were able to replicate in the host's tumor cells. These results demonstrate that the double

mutant viruses of the disclosed invention are safer than other forms of vaccinia virus, when administered to a tumor bearing host.

#### Example 7

#### Construction of a Double Mutant Vaccinia Virus Expression Vector for Expressing Cytosine Deaminase (CD)

[0109] A double mutant vaccinia virus is constructed according to the method described in Example 1 except that a gene encoding cytosine deaminase (CD) is incorporated into the shuttle vector containing the TK gene.

[0110] The CD gene containing shuttle vector is introduced into the vaccinia virus host cell for homologous recombination using a liposome mediated transfection protocol. Briefly, a 6-well tissue culture plate is seeded with  $5 \times 10^5$  CV-1 cells/well in complete DMEM-10 medium. The cells are grown to near confluency. The VGF- VSC20 virus of Example 1 is prepared and used to infect the monolayer of CV-1 cells for a period of 30 minutes. Approximately 5 minutes before the end of the infection, a liposome suspension is prepared according to the method of Whitt et al., Current Techniques in Molecular Biology, (Ed. Ausubel, et al.) Unit 9.4 (1998). In a 12x75-mm polystyrene tube is placed 1 ml of DMEM medium (Biofluids; Rockville, MD) for each well to be tested. The liposome suspension is vortexed and 15  $\mu$ l of the suspension is added to the medium. The mix is vortexed again. To this solution is added 5  $\mu$ g of shuttle vector DNA, followed by gentle mixing. Following preparation of the liposome solution, the virus inoculum is aspirated from the CV-1 cells, and the DNA/liposome complex is added directly to the cells. The cells are incubated 4 hours.

[0111] Utilizing the method of homologous recombination, the CD gene under the control of the vaccinia virus synthetic early/late promoter (See TABLE 1) is introduced into the vaccinia virus genome of VSC20. Since the VSC20 virus already contains a mutation in the VGF gene, it has a negative VGF phenotype. After the introduction of the CD gene into the TK gene locus by homologous recombination from the shuttle vector, the resulting double mutant vaccinia virus possesses a negative phenotype for VGF, a negative phenotype for TK, and it will cause cells infected with this virus to express the CD gene.

[0112] Virus particles containing the CD gene are identified using positive selection pressure and repeated rounds of plaque purification described above.

#### Example 8

##### In Vivo Antitumor Response of Double Mutant Vaccinia Virus

##### Following Systemic Injection

[0113] This Example discusses data gathered from a tumor bearing host injected systemically with the double mutant virus. Approximately  $10^5$  MC-38 (murine colon cancer) cells in 100 $\mu$ l of DMEM were injected subcutaneously into the right flanks of 6-week old female athymic (C57Bl6) mice. When the tumors reached 75 to 125 mm<sup>3</sup> in volume,  $10^9$  pfu of VVDDEGFP or HBSS (Hanks Balanced Salt Solution, Biofluids, Chicago, IL) control was injected intraperitoneally in 2ml of HBSS/0.1%BSA (Calbiochem, La Jolla, CA). Tumor width and length was measured twice weekly by a blinded investigator. Tumor volume was calculated as  $[(width)^2 \times length]0.52$ . Tumor volume was graphed as a function of time. The data from this experiment is shown in FIGURE 6.

[0114] The data shown in FIGURE 6 demonstrates animals administered a double mutant vaccinia virus of the disclosed invention experience little or no tumor volume increase, as opposed to animals that received medium alone. These results show that administration of a double mutant vaccinia virus of the disclosed invention to a tumor bearing-host animal can significantly impact tumor growth in that host.

#### Example 9

##### In Vivo Use of Double Mutant Vaccinia Virus Carrying HER2 Gene In Combination With Monoclonal Antibody Therapy

[0115] In this Example, a tumor-bearing host is infected with a double mutant vaccinia virus that carrying the gene for the HER2 protein. The virus infected tumors express the HER2 protein. Once expression of the HER2 protein has occurred, the host is provided an efficacious dose of Herceptin (Trastuzumab) (Genentech, South San Francisco, CA). Herceptin is an IgG 1 kappa monoclonal antibody that contains human frame-work regions with the complementarity-determining regions of a murine antibody (4D5) that binds to the extracellular domain of the human epidermal growth factor receptor2 protein, HER2. This receptor can be expressed by the virus and thus improve the effect of Herceptin.

[0116] Cells infected with the virus will be killed either through the replication of the virus or through the action of the monoclonal antibody. Further, radio-labeled or chemo-labeled antibodies are also used in this treatment regime. The labeled antibodies will kill HER2 expressing cells and by-stander non-infected cells, as well. This treatment results in the elimination of tumor cells.

#### Example 10

##### Use of Double Mutant Vaccinia Virus Carrying Somatostatin Gene

##### To Facilitate Tumor Imaging

[0117] A patient with a known carcinoid tumor is infected with a double deleted virus carrying the somatostatin receptor. The virus infects tumor cells and expresses the receptor. The patient then receives an intravenous injection of 5.5 millicuries of Indium 111 labeled Octreotide to image somatostatin receptor-expressing cells.

[0118] Whole body images demonstrate relatively normal uptake in liver, spleen, kidney and bladder. There is however a diffuse mid-abdominal small bowel uptake which is beyond the usual amount of activity seen on both the 4 hour and 24 hour images. This amount of activity is not normal and is most likely small bowel involvement with carcinoid. SPECT activity of the liver demonstrates multiple focal areas of increased uptake seen with multiple metastatic carcinoid deposits in the liver. The results of this experiment indicate the utility of this procedure with which to image tumors in a subject.

#### Example 11

##### Use of Double Mutant Vaccinia Virus Carrying Transferrin Receptor Gene

##### To Facilitate Tumor Imaging

[0119] Another example of tumor imaging exploits work recently published (Nat. Med. Mar;6(3):351-5(2000)) using the transferrin receptor to enhance uptake of an iron-based contrast material used for MRI imaging. A double mutant vaccinia virus carrying the transferrin receptor (ETR) gene is engineered to express the ETR gene as a reporter that shuttles targeted superparamagnetic nanoparticles into virus infected tumor cells. The internalizing receptor used is a human transferrin receptor that lacks the iron-regulatory region and mRNA destabilization motifs in the 3' untranslated region and therefore constitutively overexpresses high levels of the receptor protein in the tumor cell. These

features result in approximately a 500% increase in holo-transferrin bound by stably transfected cells in culture. The receptor-targeted MR reporter consists of 3-nm monocrystalline iron oxide nanoparticles (MION), sterically protected by a layer of low-molecular-weight dextran to which human holo-transferrin (Tf) is covalently conjugated (Tf-MION). Several synergistic effects theoretically contribute to visualizing transgene expression *in vivo*.

#### Example 12

##### Inducing Antibodies Against an Antigen Expressed by a Vaccinia Virus Double Mutant Expression Vector

[0120] T cells and B cells both comprise cell surface proteins which can be utilized as "markers" for differentiation and identification. One such human B cell marker is the human B lymphocyte-restricted differentiation antigen Bp35, referred to as "CD20." CD20 is expressed during early pre-B cell development and remains until plasma cell differentiation. Specifically, the CD20 molecule may regulate a step in the activation process which is required for cell cycle initiation and differentiation and is usually expressed at very high levels on neoplastic ("tumor") B cells. CD20, by definition, is present on both "normal" B cells as well as "malignant" B cells, ie, those B cells whose unabated proliferation can lead to B cell lymphoma. Thus, the CD20 surface antigen has the potential of serving as a candidate for "targeting" of B cell lymphomas.

[0121] Antibodies raised in an individual suffering from a B cell lymphoma are efficacious as a therapeutic modality. Accordingly, the double mutant vaccinia virus expression vector of the present invention can be used to express a B cell lymphoma specific antigen in a subject suffering from B cell lymphoma.

[0122] To treat an individual presenting a B cell lymphoma, a double mutant vaccinia virus expression vector is constructed according to the methods described in Examples 1 and 2. Incorporated into the double mutant expression vector is an antigenic portion of the CD20 surface antigen. An infectious dose of the double mutant vaccinia virus expression vector encoding the CD20 antigen is provided to a subject, and thus the cells of that subject. The mutant vaccinia virus construct enters into an appropriate host cell and



begins to express the CD20 antigen. The immune system of the treated subject then responds to the expression of the antigen and mounts an antibody response to the CD 20 antigen.

#### Example 13

##### Inducing A Cellular Response Against an Antigen Expressed by a Vaccinia Virus Double Mutant Expression Vector

[0123] The human immune system can be divided generally into a humoral category and a cellular category. Antibodies and the complement system comprise the humoral system while cytotoxic T cells, NK cells, macrophages, and other cell types comprise the cellular category. The vaccinia virus expression vectors of the present invention can be used to raise a cellular immune responses from a treated individual.

[0124] To induce a cellular immune response in an individual, a double mutant vaccinia virus expression vector is constructed according to the methods described in Examples 1 and 2. It is known that the human immunodeficiency virus (HIV) *gag* protein can induce cellular immune response when used to immunize a mammalian subject. Accordingly, an antigenic portion of the HIV *gag* protein is incorporated into the double mutant expression vector. An infectious dose of the double mutant vaccinia virus expression vector encoding the HIV *gag* protein is provided to a subject, and thus the cells of that subject. The mutant vaccinia virus construct enters into an appropriate host cell and begins to express the HIV *gag* protein. The immune system of the treated subject then responds to the expression of the antigen and generates a cellular response to the HIV *gag* protein.

#### Example 14

##### Inducing Cell Death by Infecting a Tumor Cell with a Vaccinia Virus Double Mutant Expression Vector Containing a Suicide Gene

[0125] Murine colon adenocarcinoma cells (MC38 cells) are grown to semi confluency and then infected with the double mutant vaccinia virus expression vector described in Example 7 that expresses the cytosine deaminase (CD) gene. A control culture of MC38 cells is also established in which MC38 cells are grown and infected with a double mutant vaccinia virus that does not contain the CD gene. Following infection of both cell cultures a dose of 5-fluorocytosine (5-FC) is added to both cultures. The dose of 5-FC is of

